

**ROBUST SUMMARY  
OF INFORMATION ON**

**Substance Group**

**Kerosene/Jet Fuel**

**Summary prepared by**

American Petroleum Institute  
Consortium Registration # 1100997

**Creation date:**

May 24, 2003

**Printing date:**

September 21, 2010

**Date of last Update:**

**September 21, 2010**

**Number of pages:**

**119**

NB. Reliability of data included in this summary has been assessed using the approach described by Klimisch et al.

Klimisch, H. J., Andreae, M. and Tillman, U, (1997)

A systematic approach for evaluating the quality of experimental toxicological and ecotoxicological data.

Regulatory Toxicology and Pharmacology 25, 1-5.

# 1. General Information

Id Kerosene  
Date 9/21/2010

## 1.1.1 GENERAL SUBSTANCE INFORMATION

**Substance type** : Petroleum product  
**Physical status** : Liquid

**Remark** : Kerosene is the name for the lighter end of a group of petroleum streams known as the middle distillates.

Kerosene may be obtained either from the distillation of crude oil under atmospheric pressure (straight-run kerosene) or from catalytic, thermal or steam cracking of heavier petroleum streams (cracked kerosene). The kerosenes, are further treated by a variety of processes (including hydrogenation) to remove or reduce the level of sulfur, nitrogen or olefinic materials.

The precise composition of any particular kerosene will depend on the crude oil from which it was derived and on the refinery processes used for its production. Irrespective of this, kerosenes consist predominantly of C9 to C16 hydrocarbons and boil in the range 145 to 300 °C. The major components of kerosenes are branched and straight chain paraffins and naphthenes and these normally account for 70% of the material. Aromatic hydrocarbon, mainly alkyl benzenes and alkylnaphthalenes do not normally exceed 25 % of kerosene streams. Olefins do not normally account for more than 5% of the kerosenes.

Several mammalian toxicity studies have been carried out on two samples of kerosene and the properties of these are shown below. Sample 83-09 is a straight-run kerosene and 81-07 is a hydrodesulfurized kerosene.

<b>Property</b>	<b>Sample API 83-09</b>	<b>Sample API 81-07</b>
CAS number	8008-20-6	64742-81-0
Pour point (°C)	-49	
Density (kg/dm <sup>3</sup> )@ 15 °C	0.81	0.82
Reid vapor pressure @ 37.8 °C (hPa)	14	
Flash point (closed cup) °C	62	60
Kinematic viscosity at 20 °C (mm <sup>2</sup> /sec)	1.5-2.5	1.1-2.5
Gravity (°API)	43.0	41.9
Sulfur (wt %)	0.47	0.07
Nitrogen (ppm)	5.4	-
Flash point (°F)	144	140
Distillation (°F)		
IBP	238	362
10%	327	392
50%	405	434
90%	475	488
95%	490	506
EP	520	535
Paraffins (%)	-	47
Saturates (%)	82	-
Olefins (%)	2.5	1
Naphthenes (%)	-	35
Aromatics (%)	15.5	18

# 1. General Information

Id Kerosene  
Date 9/21/2010

The major use of kerosenes is as the primary ingredient in a variety of jet fuels. Kerosene is also used as diesel fuel (No. 1), domestic heating fuels (Fuel oil No. 1) and as a solvent, although this latter use is a minor one. Both diesel fuel and home heating oil No. 1 are essentially equivalent to kerosene. Specifications for both middle distillate heating fuels and transportation fuels are similar; as a consequence, it is often possible for refiners to satisfy the performance requirements of both applications with the same process stream or blend of process streams. The final products are essentially kerosene that contains additives which are specific for the intended use. Otherwise they are all virtually indistinguishable on the basis of their gross physical or chemical properties.

Characteristics of the various jet fuels are:

Jet A-1	Kerosene type used in civil aircraft, aromatic hydrocarbon content 25% (v/v) maximum, Freezing point -47°C max.
Jet A	as for Jet A-1, but with freezing point -40°C max.
AVCAT/JP-5	high flash point kerosene type used in naval aircraft, aromatic hydrocarbons 25% (v/v) max.
AVTUR/JP-8	kerosene type used in military aircraft, aromatic hydrocarbons 25 % (v/v) max.

For most of the mammalian toxicology endpoints, information has been derived on a wide range of kerosene streams and jet fuels. For simplicity, this robust summary contains detailed information on a single API sample for each endpoint and if data were available on other samples for the same endpoint they have been summarized in tabular form in the relevant sections or discussed in detail when appropriate.

## 1.13 REVIEWS

**Memo** : IARC

**Remark** : IARC reviewed the data on the carcinogenicity of jet fuel and concluded:

There is inadequate evidence for the carcinogenicity in humans of jet fuel.

There is inadequate evidence for the carcinogenicity in experimental animals of jet fuel.

The IARC overall evaluation was that  
Jet fuel is not classifiable as to its carcinogenicity to humans.

(55)

**Memo** : CONCAWE

**Remark** : CONCAWE summarized the available health, safety and environmental data available on kerosene and jet fuels

(36)

**Memo** : ASTDR

**Remark** : A Toxicological profile for JP-5 and JP-8 jet fuel was prepared by the Agency for Toxic Substances and Disease Registry

(28)

# 1. General Information

**Id** Kerosene  
**Date** 9/21/2010

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

#### 2.1 MELTING POINT

**Method** : ASTM D97  
**GLP** : No data  
**Test substance** : Kerosene/Jet Fuels various

**Remark** : By definition, melting point is the temperature at which a solid becomes a liquid at normal atmospheric pressure. For complex mixtures like petroleum products, melting point may be characterized by a range of temperatures reflecting the melting points of the individual components. To better describe phase or flow characteristics of petroleum products, the pour point is routinely used. The pour point is the lowest temperature at which movement of the test specimen is observed under prescribed conditions of the test (ASTM 1999). The pour point methodology also measures a "no-flow" point, defined as the temperature of the test specimen at which a wax crystal structure and/or viscosity increase such that movement of the surface of the test specimen is impeded under the conditions of the test (ASTM 1999). Because not all petroleum products contain wax in their composition, the pour point determination encompasses change in physical state (i.e., crystal formation) and/or viscosity property. Given the pour point values given above, these products are liquid at ambient temperatures.

The pour point values for the products and streams given above are expected to represent the pour point values for the entire Kerosenes/Jet Fuel HPV category. This is expected because:

- 1) the products within this category consist of the same types of petroleum hydrocarbons (paraffinic, naphthenic, olefinic and aromatic),
- 2) the proportions of those hydrocarbon types are similar among the products/streams,
- 3) the hydrocarbons have a relatively similar and narrow range of molecular weights (carbon atoms of C9 - C16), and
- 4) the products and streams have similar distillation ranges (approximately 125 to 300 °C).

**Result** : Pour point: See following Table and Remarks Section  
Decomposition: N/A  
Sublimation: N/A

<b>Product Type</b>	<b>Pour Point (°C)</b>	<b>Reference</b>
CAS No. 8008-20-6 (Straight-run kerosene)	-55	API 1987
Jet A/Jet A-1 (aviation turbine fuel, kerosene type)	< -47	Jokuty, et al. 2002
Fuel Oil No. 1 (JP-8; kerosene)	-50	Jokuty, et al. 2002
Fuel Oil No. 1 (JP-5; heavy kerosene)	< -48	Jokuty, et al. 2002

**Reliability** : (2) valid with restrictions  
Results of standard method testing were reported in a reliable reference database.

(22) (29) (56)

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

#### 2.2 BOILING POINT

**Method** : ASTM D86  
**GLP** : No data  
**Test substance** : Kerosene/Jet Fuels various

**Remark** : The boiling (distillation) ranges given below are expected to cover other streams and/or products in the Kerosene/Jet Fuel HPV category. This is expected due to their similar hydrocarbon content representing paraffinic, naphthenic, olefinic and aromatic constituents and their proportions within the substances in this category. Data presented below for streams not included in the Kerosene/Jet Fuel HPV category (e.g., CAS Nos. 91770-15-9 and 101316-80-7) provide supporting information to illustrate the overall similarity in boiling range for products within this category.

The one stream within the Kerosene/Jet Fuel HPV category having a larger range of molecular weight hydrocarbons than other members of the category (i.e., CAS No. 68477-58-7 with C5 to C18 carbon atoms) would be expected to have a boiling (distillation) range wider than those characterized by the cited data.

**Result** : Boiling point (°C): See following Table  
Pressure: N/A  
Pressure unit: N/A  
Decomposition: N/A

<b>Boiling Range °C</b>	<b>Method</b>	<b>Ref.</b>
Jet A/Jet A-1 145 - 300	ASTM D86	Jokuty et al. 2002
Kerosene, straight-run 125 - 292	API 83-09; (CAS No. 8006-20-6) ASTM D86	API 1987
Kerosene, hydrodesulfurized 175 - 284	API 81-08, (CAS No. 64742-81-0) ASTM D86	API 1987
Kerosene, sweetened (CAS No. 91770-15-9) 152 - 257	ASTM D86	CONCAWE 94/106
Kerosene, hydrodesulfurized (CAS No. 64742-81-0) 156 - 255	ASTM D86	CONCAWE 94/106
Kerosene, hydrocracked heavy aromatic (CAS No. 101316-80-7) 187 - 288	ASTM D86	CONCAWE 96/55

**Reliability** : (2) valid with restrictions  
Results of standard method testing were reported in reliable review dossiers and a reference database.

(22) (39) (56)

#### 2.4 VAPOUR PRESSURE

**Method** : Measured: ASTM 323

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

**Test substance** : Kerosene/Jet Fuels various

**Remark** : Kerosenes and jet fuels consist of complex mixtures of various hydrocarbon compounds with the majority of the structures represented by saturated hydrocarbons (e.g., normal and branched paraffins and naphthenes) with lesser amounts of aromatic compounds. Molecular weights of those constituents are represented by components containing between nine and sixteen carbon atoms (CONCAWE 1995).

**Result** : Although individual hydrocarbon constituents in kerosenes exert their own vapor pressures, the vapor pressure of the mixture is the sum of the individual partial pressures of the components (Dalton's Law). The values given below are considered to be representative of the general category of kerosenes and jet fuels and the petroleum refining streams that produce them.

Vapor pressure: See table below:  
Temperature: Not given  
Decomposition: Not given

**Vapor Pressures**

kPa:	Temp (°C)	Method	Ref.
JP-5 0.3 - 3.5	21	Not Given	Air Force (1989)
JP-8 0.3 - 3.5	21	Not Given	Air Force (1989)
Jet A/Jet A-1 > 1	37.8	ASTM D323	Jokuty et al.(2002)
Straight-run kerosene 1.4	37.8	CAS No. 8008-20-6 ASTM D323	API (1987)

**Reliability** : (2) valid with restrictions  
Results of standard method testing were reported in reliable review dossiers and a reference database.

(1) (22) (30) (36) (56)

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

#### 2.5 PARTITION COEFFICIENT

##### Category: KEROSENE/JET FUELS

Category Chemical: kerosene, various CAS RN

Test Substance: kerosene, various CAS RN

Test Substance  
Purity/Composition and  
Other Test Substance  
Comments:

Category Chemical Result  
Type: Estimated by calculation

Test Substance Result  
Type: Estimated

#### RESULTS

Partition Coefficient Input type: Value or Range?  
Values are given for individual gas oil constituent hydrocarbons.

Partition Coefficient Range : Log  $K_{ow}$ : 3.3 - >6 @ Temperature: 25°C

Representative gas oil constituent	C-num	PC (Log Kow)
<b>Paraffins (n-, iso-)</b>		
n-nonane	9	4.8
iso-nonane	9	4.7
n-hexadecane	16	8.2
iso-hexadecane	16	8.1
<b>Naphthenes</b>		
propylcyclohexane	9	4.6
bicyclo[4.3.0]nonane	9	3.7
n-decylcyclohexane	16	8.0
2-hexyl[4.4.0]bicyclodecane	16	7.1
<b>Olefins</b>		
1-nonene	9	5.2*
1-hexadecene	16	8.1
<b>Aromatic</b>		
n-propylbenzene	9	3.7*
naphthalene	10	3.3*
n-decylbenzene	16	7.4*
3-hexylnaphthalene	16	6.2

Calculated Log Kow were performed using EPI-Suite V3.10, KOWWIN V1.66 (EPA 2000). Values in the above table that were cited in the KOWWIN experimental database are denoted by "\*".

#### Results Remarks:

Calculated partition coefficients (log Kow) are given for various paraffinic, olefinic, naphthenic, and aromatic structures having carbon numbers from C9 to C16, which encompasses the principal carbon number range given for the category. The values are consistent with the range of 3.3 to >6 calculated by CONCAWE (1995). These Kow estimates cover kerosene streams and fuels.

#### STUDY/METHOD



### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

**Key Study Sponsor Indicator:**

**Year Study Performed:**

**Method/Guideline Followed:** Subroutine KOWWIN V1.66 in EPI-Suite V3.10

**Method/Guideline and Test Condition Remarks:**

**GLP:** No

**Study Reference:** U.S. EPA. 2000. EPI (Estimation Programs Interface) Suite, V3.10. U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics, Washington, DC.

#### RELIABILITY/DATA QUALITY

**Reliability:** (2) Reliable with restrictions

**Reliability Remarks:** Estimated values were derived from a recognized validated computer model.

#### 2.6.1 SOLUBILITY IN DIFFERENT MEDIA

**Category Chemical :** Kerosene/Jet Fuels

**Test Substance :** Jet A-1 fuel

**Test Substance Purity/Composition and Other Test Substance Comments:**

**Category Chemical Result Type :** Measured

**Test Substance Result Type :** Measured

#### RESULTS

**Water Solubility Indicator :**

**Water Solubility Input type:** Value

**Water Solubility Value/Range :** **Solubility:** 10.4 mg/L @ **Temperature:** 20°C

**pH Value :** **Value or Lower Range:** N/A  
**Upper Range :**

**pKa - Protein Kinase:** N/A

**pH Value at Saturation :** N/A

**Results Remarks :**

#### STUDY/METHOD

**Key Study Sponsor Indicator :**

**Year Study Performed :** 1994

**Method/Guideline Followed :** Other

**Method/Guideline and Test** Water soluble fractions were created using a ratio of fresh well water to

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

**Condition Remarks:**

oil of 40:1 (v/v) in 2-L Erlenmeyer flasks fitted with Teflon stopcocks near the bottom for drainage. The caps were lined with teflon tape and the flasks were covered with foil to minimize possible photodegradation. Water was added to the flasks along with a Teflon stir-bar followed by the oil. Flasks were capped and placed on a magnetic stirrer for three days of stirring. Stirring was adjusted to approximately one revolution per second so that no vortex was formed on the underside of the oil and no oil emulsion was created.

Samples of the water soluble fraction were collected in amber septum bottles then analyzed by purge and trap gas chromatography. Instrumentation included a Texmar Model 4000 Dynamic Headspace Concentrator coupled to a cryofocusing Texmar Model 1000 Capillary interface and a Hewlett-Packard Model 5890 gas chromatograph equipped with a flame ionization detector and a SPB-1 fused silica column (0.5e mm ID x 30 m long).

Instrument conditions:

Purge and Trap:	GC Programming:
Purge time: 15 min	40°C for 10 min
Desorb time: 6 min	8°C/min
Bake time: 25 min	225°C for 10 min

**GLP :**

Not stated

**Study Reference :**

EC (Environment Canada). 1994. The comparative toxicity of crude and refined oils to Daphnia magna. Unpublished report series EE-152, Environmental Technology Centre, Ottawa, Canada. 23 pp.

**RELIABILITY/DATA QUALITY**

**Reliability :**

2

**Reliability Remarks :**

Reliable with restrictions. Meets basic scientific principles but lacks some details concerning preparation of the water soluble fraction.

<b>Category Chemical :</b>	Kerosene/Jet Fuels
<b>Test Substance :</b>	Kerosene-type jet fuel
<b>Test Substance Purity/Composition and Other Test Substance Comments:</b>	Distillation range 200 to 300°C
<b>Category Chemical Result Type :</b>	Measured
<b>Test Substance Result Type :</b>	Measured
<b>RESULTS</b>	
<b>Water Solubility Indicator :</b>	
<b>Water Solubility Input type:</b>	Value
<b>Water Solubility Value/Range : Solubility:</b> 4.78 mg/L @ <b>Temperature:</b> 27°C	
<b>pH Value :</b>	<b>Value or Lower Range:</b> N/A <b>Upper Range :</b>
<b>pKa - Protein Kinase:</b>	N/A

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

<b>pH Value at Saturation :</b>	N/A																		
<b>Results Remarks :</b>																			
<b>STUDY/METHOD</b>																			
<b>Key Study Sponsor Indicator :</b>																			
<b>Year Study Performed :</b>	1998																		
<b>Method/Guideline Followed :</b>	Other																		
<b>Method/Guideline and Test Condition Remarks:</b>	<p>The water soluble fraction was created using a ratio of culture medium to oil of 200:1. The solution was stirred for five hours and the insoluble fraction was separated.</p> <p>Samples of the water soluble fraction were analyzed by gas chromatography/mass spectroscopy (GC-MS) using the following instruments and conditions:</p> <p>Instrument: Varian 3400CX gas chromatograph Column: OV-101 (length 30m, ID 0.25mm) Packing: DB-1 (1µm dia)</p> <table> <tr> <td>Operating conditions:</td> <td>GC programming:</td> </tr> <tr> <td>Carrier gas: nitrogen</td> <td>40°C for 2 min</td> </tr> <tr> <td>Flow rate: 3 mL/min</td> <td>increase 25°C/min for 7.2 min until</td> </tr> <tr> <td>Pressured air: 350 mL/min</td> <td>220°C</td> </tr> <tr> <td>Hydrogen: 30 mL/min</td> <td>Hold 1 min</td> </tr> <tr> <td>Nitrogen: 25 mL/min</td> <td>Increase to 300°C @ 15°C/min</td> </tr> <tr> <td>Detector: flame ionization</td> <td>Detector temp: 250°C</td> </tr> <tr> <td>Temperature: 300°C</td> <td></td> </tr> <tr> <td>Injection volume: 1 or 2 µL</td> <td></td> </tr> </table>	Operating conditions:	GC programming:	Carrier gas: nitrogen	40°C for 2 min	Flow rate: 3 mL/min	increase 25°C/min for 7.2 min until	Pressured air: 350 mL/min	220°C	Hydrogen: 30 mL/min	Hold 1 min	Nitrogen: 25 mL/min	Increase to 300°C @ 15°C/min	Detector: flame ionization	Detector temp: 250°C	Temperature: 300°C		Injection volume: 1 or 2 µL	
Operating conditions:	GC programming:																		
Carrier gas: nitrogen	40°C for 2 min																		
Flow rate: 3 mL/min	increase 25°C/min for 7.2 min until																		
Pressured air: 350 mL/min	220°C																		
Hydrogen: 30 mL/min	Hold 1 min																		
Nitrogen: 25 mL/min	Increase to 300°C @ 15°C/min																		
Detector: flame ionization	Detector temp: 250°C																		
Temperature: 300°C																			
Injection volume: 1 or 2 µL																			
<b>GLP :</b>	Not stated																		
<b>Study Reference :</b>	Rozkov, A., A. Kaard, and R. Vilu. 1998. Biodegradation of dissolved jet fuel in chemostat by a mixed bacterial culture isolated from a heavily polluted site. Biodegradation 3:363-369.																		
<b>RELIABILITY/DATA QUALITY</b>																			
<b>Reliability :</b>	2																		
<b>Reliability Remarks :</b>	Reliable with restrictions. Meets basic scientific principles but lacks some details concerning preparation of the water soluble fraction.																		
<b>Category Chemical :</b>	Kerosene/Jet Fuels																		
<b>Test Substance :</b>	JP-8 jet fuel																		
<b>Test Substance Purity/Composition and Other Test Substance Comments:</b>	Flash point range: 40.56 – 65.66°C Specific gravity range: 0.775 – 0.840																		
<b>Category Chemical Result Type :</b>	Measured																		
<b>Test Substance Result Type :</b>	Measured																		

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

#### RESULTS

<b>Water Solubility Indicator :</b>	
<b>Water Solubility Input type:</b>	Value
<b>Water Solubility Value/Range :</b>	<b>Solubility:</b> mean: 12.44 mg/L $\pm$ 0.71 @ <b>Temperature:</b> not stated
<b>pH Value :</b>	<b>Value or Lower Range:</b> N/A <b>Upper Range :</b>
<b>pKa - Protein Kinase:</b>	N/A
<b>pH Value at Saturation :</b>	N/A
<b>Results Remarks :</b>	The mean concentration of the water soluble fraction for 69 replicate measurements was 10.98 mg/L ( $\pm$ 0.94). Measurements included in the average covered different days following replenishment of the solubilizer. The highest concentration typically was the first measurement taken after replenishment. This occurred at 18 h and averaged 12.44 $\pm$ 0.71 mg/L. Concentrations in the water soluble fraction declined with time following replenishment. Concentrations taken after 7 days averaged 10.67 $\pm$ 0.71 mg/L.

#### STUDY/METHOD

<b>Key Study Sponsor Indicator :</b>	
<b>Year Study Performed :</b>	1983
<b>Method/Guideline Followed :</b>	Other
<b>Method/Guideline and Test Condition Remarks:</b>	<p>The water soluble fraction was created using solubilizer device described by Krugel et al. (1978). This consisted of five glass columns connected in series and each containing 1 L fuel and 0.7 L of water. Tap water flowed down through each column and from the last column entered a 9-L glass carboy which served as a filter and separation chamber. The water soluble fraction was collected from the glass carboy.</p> <p>Samples of the water soluble fraction were analyzed by gas chromatography (GC) using the following methods, instruments, and conditions:</p> <p>The water soluble fraction (300 mL) was extracted with pentane and reduced by rotary evaporation to 3 mL. A 2-<math>\mu</math>L aliquot t was injected into a gas chromatograph (Fisher Model 4800 with dual FID and two 6.1m stainless-steel columns of 10% WE 30 on 80/100 Chrom W.</p> <p>Operating conditions: Injector temp: 270°C Detector temp: 200°C Column temp: 160°C with isothermal operation The concentration of hydrocarbons in the water soluble fraction was reported as the sum of the total peak area of all components.</p>
<b>GLP :</b>	Not stated
<b>Study Reference :</b>	Klein, S.A., and D. Jenkins. 1983. The toxicity of jet fuels to fish – II. The toxicity of JP-8 to flagfish ( <i>Jordanella florida</i> ) and rainbow trout ( <i>Salmo gairdneri</i> ) and golden shiners ( <i>Notemigonus chysoleucas</i> ). Wat. Res. 17(10):1213-1220.

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

#### RELIABILITY/DATA QUALITY

<b>Reliability :</b>	2
<b>Reliability Remarks :</b>	Reliable with restrictions. Meets basic scientific principles but lacks some details concerning preparation of the water soluble fraction.

#### WATER SOLUBILITY

<b>Category Chemical :</b>	Kerosene/Jet Fuels
<b>Test Substance :</b>	No. 1 fuel oil
<b>Test Substance Purity/Composition and Other Test Substance Comments:</b>	
<b>Category Chemical Result Type :</b>	Measured
<b>Test Substance Result Type :</b>	Measured

#### RESULTS

<b>Water Solubility Indicator :</b>																	
<b>Water Solubility Input type:</b>	Value																
<b>Water Solubility Value/Range : Solubility:</b> 5.16 – 5.52 mg/L @ <b>Temperature:</b> not stated																	
<b>pH Value :</b>	<b>Value or Lower Range:</b> N/A <b>Upper Range :</b>																
<b>pKa - Protein Kinase:</b>	N/A																
<b>pH Value at Saturation :</b>	N/A																
<b>Results Remarks :</b>	<p>The 100% water soluble fractions (WSF) were diluted to 50% prior to measurement. Three measurements were taken using the combined analysis technique. Concentrations were multiplied by 2 to obtain the calculated concentrations in the 100% soluble fraction sample.</p> <table border="1"> <thead> <tr> <th>Trial#</th> <th>Headspace analysis, mg/L</th> <th>Micro-extraction analysis, mg/L</th> <th>Calculated Conc. in 100% WSF, mg/L</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>0.25</td> <td>2.33</td> <td>5.16</td> </tr> <tr> <td>2</td> <td>0.25</td> <td>2.36</td> <td>5.22</td> </tr> <tr> <td>3</td> <td>0.25</td> <td>2.51</td> <td>5.52</td> </tr> </tbody> </table>	Trial#	Headspace analysis, mg/L	Micro-extraction analysis, mg/L	Calculated Conc. in 100% WSF, mg/L	1	0.25	2.33	5.16	2	0.25	2.36	5.22	3	0.25	2.51	5.52
Trial#	Headspace analysis, mg/L	Micro-extraction analysis, mg/L	Calculated Conc. in 100% WSF, mg/L														
1	0.25	2.33	5.16														
2	0.25	2.36	5.22														
3	0.25	2.51	5.52														

#### STUDY/METHOD

<b>Key Study Sponsor Indicator :</b>	
<b>Year Study Performed :</b>	1987
<b>Method/Guideline Followed :</b>	Other
<b>Method/Guideline and Test Condition Remarks:</b>	<p>Water soluble fractions were generated by mixing water and oil at a ratio of 80:1 water/oil. Mixtures were placed in Mariotte bottles that were completely filled without headspace. The solutions were mixed for 2 hours after which they were allowed to settle for 72 hours and separate into aqueous and oil phases.</p> <p>Aliquots of the water soluble fractions were analyzed by two methods,</p>

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

	<p>headspace analysis and micro-extraction analysis. The headspace analysis was used to measure the extremely volatile low-boiling hydrocarbons with boiling points up to 115°C. This was followed by solvent micro-extraction, which measured the hydrocarbons boiling between 115 and 270°C. The sum of the two analyses was taken as the total hydrocarbon in the water soluble fraction. In both measurements, hydrocarbons were measured via gas chromatography (GC) with flame ionization detector (FID).</p> <p>In the headspace method, a 25-mL water sample was equilibrated with nitrogen gas (25 mL) in a gas-tight syringe. The gas phase was then injected into a HP 5750 gas chromatograph fitted with a 3-m column of 5% Dexsil 300. Integrated peak areas were converted to weights of hydrocarbons using response factors established from authentic standard solutions of hydrocarbons.</p> <p>In the micro-extraction method, 90-mL aliquots of water were extracted into 250 µL hexane in a microextraction flask. The hexane layer was analyzed by GC using a Varian Vista 6000 fitted with a 15m DB5 wide bore capillary column with splitless injection technique.</p> <p>These two methods covered the measurement of hydrocarbons having carbon numbers from C1 (methane) to C12 (2-methyl-naphthalene).</p>
<b>GLP :</b>	Not stated
<b>Study Reference :</b>	Lockhart, W.L., R.W.Danell, and D.A.J. Murray. 1987. Acute toxicity bioassays with petroleum products: Influence of exposure conditions. Pages 335-344 <u>In</u> : Oil in Freshwater: Chemistry, Biology, Countermeasure Technology. Vandermeulen and Hrudey, eds. Pergamon Press, New York.
<b>RELIABILITY/DATA QUALITY</b>	
<b>Reliability :</b>	2
<b>Reliability Remarks :</b>	Reliable with restrictions. The analysis technique did not include the hydrocarbons from C13 to C16, which exist in members of the kerosene/jet fuel category. Therefore, reported concentrations in the water soluble fraction may not include the total hydrocarbons potentially present.

#### 2.6.2 ADDITIONAL REMARKS

- Memo Remark** : Water solubility of kerosene/jet fuels  
: ATSDR (1998) cited a water solubility value of approximately 5 mg/l at 20 °C for JP-5/JP-8 (kerosene) that had been reported by the Air Force (1989).
- Source** : Air Force (1989)  
The installation restoration program toxicology guide. Oak Ridge, TN: Biomedical and Environmental Information Analysis. Vol. 4. [as cited in ATSDR (Agency for Toxic Substances and Disease Registry). 1998. Toxicological Profile for JP-5 and JP-8. U.S. Department of Health and Human Services, Public Health Service, Atlanta, Georgia].
- Reliability** : (4) not assignable
- Memo** : Water solubility of kerosene/jet fuels

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

#### 3.1.1 PHOTODEGRADATION

**Category:** KEROSENE/JET FUELS

**Category Chemical :** kerosene, various CAS RN

**Test Substance :** kerosene, various CAS RN

**Test Substance Purity/Composition and Other Test Substance Comments:**

**Category Chemical Result Type :** Estimated by calculation

**Test Substance Result Type :** Estimated

#### RESULTS

#### Photodegradation Result Description:

Direct photolysis is not expected to be a major degradation pathway for most of the components in kerosenes/jet fuels. Chemicals having the potential to photolyze have UV/visible absorption maxima in the range of 290 to 800 nm. Most hydrocarbon constituents in the kerosene/jet fuel category do not show absorbance within the 290-800 range. However, where unsaturated hydrocarbons, notably polyaromatic hydrocarbons are present in, or near the surface of water, degradation by reaction with sunlight in the presence of oxygen can be a removal process for those compounds (CONCAWE 2001).

Constituents of kerosenes that volatilize to the troposphere have the potential to undergo gas-phase oxidation reactions with photochemically produced hydroxyl radicals (OH), other oxygen containing radicals (e.g., NO<sub>3</sub>) and ozone (O<sub>3</sub>). Atmospheric oxidation as a result of these types of reactions is not direct photochemical degradation but indirect photodegradation (Schwarzenbach et al, 2003).

**Photodegradation Input type :** Value or Range?  
Values given for individual hydrocarbon constituents in kerosenes/jet fuels.

#### Photodegradation Range :

#### Half Life :

Representative gas oil constituent	C-num	Half-life, days
<b>Paraffins (n-, iso-)</b>		
n-nonane	9	1.1
iso-nonane	9	1.1
n-hexadecane	16	0.5
iso-hexadecane	16	0.5
<b>Naphthenes</b>		
n-propylcyclohexane	9	0.8
bicyclo[4.3.0]nonane	9	0.8
n-decylcyclohexane	16	0.5
2-hexyl[4.4.0]bicyclodecane	16	0.4
<b>Olefins</b>		

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

	1-nonene	9	0.3
	1-hexadecene	16	0.2
<b>Aromatic</b>	propylbenzene	9	1.5
	naphthalene	10	0.5
	n-decylbenzene	16	0.6
	3-hexylnaphthalene	16	0.2

**Rate Constant :**

**Photo Medium :**

**Temperature :** 25°C

**Sensitizer :** Hydroxyl radical (OH·)

**Sensitizer Concentration and Units :**  $1.5 \times 10^6$  OH·/cm<sup>3</sup>

**Light Source :** N/A

**Light Source Spectrum :** N/A

**UV/VIS Absorption Spectrum :** N/A

**Quantum Yield :** N/A

**Breakdown Products Description :** N/A

**Results Remarks :**

It is predicted from indirect photolysis modeling of C9 and C16 paraffinic, naphthenic, olefinic, and aromatic hydrocarbon compounds that volatile components in kerosenes/jet fuels will undergo atmospheric oxidation and not persist in the environment. Based on the modeled half-lives of component hydrocarbon structures, kerosenes containing primarily aromatic or saturated fractions are not expected to show significant differences in their photodegradation characteristics.

#### STUDY/METHOD

**Key Study Sponsor Indicator :** Key

**Year Study Performed :**

**Method/Guideline Followed :** Subroutine AOPWIN V1.90 in EPI-Suite V3.10

**Deviations from Method/Guideline :**

**Method/Guideline Description :**

The potential to undergo indirect photodegradation was estimated using the atmospheric oxidation potential (AOP) model subroutine (AOPWIN V1.90) of the EPI-Suite™ computer models (EPA, 2000). This model calculates a chemical half-life and an overall OH· radical reaction rate constant based on a 12-hour day and a given OH· radical concentration. This program also estimates the reaction rates and half-lives for the reaction of olefins with O<sub>3</sub>, but as described by Atkinson (1990), these rates tend to be substantially less than for those for the OH· radical. For this reason, only the half-lives for the reaction with the OH· radical are reported for the series of olefinic hydrocarbons selected for the AOP model.



### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

#### Method/Guideline and Test Condition Remarks :

GLP : No

U.S. EPA. 2000. EPI (Estimation Programs Interface) Suite, V3.10. U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics, Washington, DC.

CONCAWE (2001) Environmental classification of petroleum substances - Summary data and rationale CONCAWE, Brussels, Belgium

Study Reference : Atkinson, R. 1990. Gas-phase tropospheric chemistry of organic compounds: A review. Atmos. Environ. 24A(1):1-41.

Schwarzenbach, R.P., Gschwend, P.M., and Imboden, D.M., eds. 2003. Chapter 16: Indirect Photolysis: Reactions with Photooxidants in Natural Waters and in the Atmosphere. In: Environmental Organic Chemistry, 2<sup>nd</sup> Edition. John Wiley and Sons, Inc.

#### RELIABILITY/DATA QUALITY

Reliability : (2) Reliable with restrictions

Reliability Remarks : Estimated values were derived from a recognized validated computer model.

#### 3.1.2 STABILITY IN WATER

Test substance : Kerosene/Jet Fuels various

Remark : Hydrolysis of an organic chemical is the transformation process in which a water molecule or hydroxide ion reacts to form a new carbon-oxygen bond. Chemicals that have a potential to hydrolyze include alkyl halides, amides, carbamates, carboxylic acid esters and lactones, epoxides, phosphate esters, and sulfonic acid esters. The chemical components that comprise the gas oil category are hydrocarbons that are not subject to hydrolysis because they lack functional groups that hydrolyze.

Result : Nominal: N/A  
Measured: N/A  
Half-life T1/2: N/A  
Degradation %: N/A  
At specified pH: N/A  
Breakdown Products: N/A

Reliability : (1) valid without restriction

(54)

#### 3.3.1 TRANSPORT BETWEEN ENVIRONMENTAL COMPARTMENTS

Category Name: KEROSENE/JET FUELS

Category Chemical : kerosene, various CAS RN

Test Substance : kerosene, various CAS RN

Test Substance  
Purity/Composition

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

and  
Other Test Substance  
Comments:

Category Chemical Result Type : Estimated by calculation

Test Substance Result Type : Estimated

#### RESULTS

Fugacity/Distribution  
Result Description:

Kerosenes are substances of variable composition containing hydrocarbons with carbon numbers typically from C9 to C16. Multimedia distribution was calculated for low and high molecular weight hydrocarbon compounds (i.e., C9 and C16 hydrocarbons) representing the principal isomeric structures in kerosene (e.g., paraffins, olefins, naphthenes, and aromatics). This provided an understanding of the distribution spectrum that might be expected for constituents at each boundary of the distribution. Hydrocarbons intermediate in carbon numbers would be expected to show distribution characteristics within the range provided by the estimates for the C9 and C16 isomeric compounds.

Test Results :

Compound/Compartment	Percent Distribution					
	Air	Water	Soil	Sed	Susp. Sed.	Biota
<b>Paraffins</b>						
n-nonane	99	<0.1	1	<0.1	<0.1	<0.1
iso-nonane	99	<0.1	0.5	<0.1	<0.1	<0.1
n-hexadecane	47	<0.1	52	1	<0.1	<0.1
iso-hexadecane	60	<0.1	36	0.9	<0.1	<0.1
<b>Naphthenes</b>						
propylcyclohexane	99	<0.1	0.9	<0.1	<0.1	<0.1
bicyclo[4.3.0]nonane	99	0.2	1	<0.1	<0.1	<0.1
n-decylcyclohexane	4.7	<0.1	93	2	<0.1	<0.1
2-hexyl[4.4.0]bicyclodecane	34	<0.1	65	1	<0.1	<0.1
<b>Olefins</b>						
1-nonene	99	<0.1	0.7	<0.1	<0.1	<0.1
1-hexadecene	10	<0.1	88	2	<0.1	<0.1
<b>Aromatics</b>						
n-propylbenzene	97	1	2	<0.1	<0.1	<0.1
naphthalene	77	8	15	0.3	<0.1	<0.1
n-decylbenzene	13	<0.1	85	2	<0.1	<0.1
3-hexylnaphthalene	1.1	<0.1	97	2	<0.1	<0.1

Temperature : 20°C

Level of Multi-media Model : 1

### 3. Environmental Fate and Pathways

Id Kerosene

Date 9/21/2010

**Model Input (Water Solubility:)**

	Water Solubility mg/L	Reference
<b>Paraffins (n-, iso-)</b>		
n-nonane	0.12	(1)
iso-nonane	0.16	(1)
n-hexadecane	0.0009*	(2)
iso-hexadecane	0.0012	(2)
<b>Naphthenes</b>		
propylcyclohexane	3.4	(2)
bicyclo[4.3.0]nonane	19	(2)
n-decylcyclohexane	0.0013	(2)
2-hexyl[4.4.0]bicyclodecane	0.0089	(2)
<b>Olefins</b>		
1-nonene	1.1*	(2)
1-hexadecene	0.0012	(2)
<b>Aromatics</b>		
n-propylbenzene	52.2*	(2)
naphthalene	31*	(2)
n-decylbenzene	0.0024*	(2)
3-hexylnaphthalene	0.20	(2)
(1) Yaws, et al., 1993 (2) US EPA, 2000 (WSKOWWIN, EPI-Suite V3.10) Note: Model input data signified by "*" indicates the value was cited by EPI-Suite™ as being from the experimental database (EPA 2000).		

**Model Input (Vapor Pressure:)**

	Vapor Pressure Pa	Reference
<b>Paraffins (n-, iso-)</b>		
n-nonane	593*	(1)
iso-nonane	828*	(1)
n-hexadecane	0.191*	(1)
iso-hexadecane	4.1	(1)
<b>Naphthenes</b>		
propylcyclohexane	558*	(1)
bicyclo[4.3.0]nonane	320	(1)
n-decylcyclohexane	0.137*	(1)
2-hexyl[4.4.0]bicyclodecane	1.1	(1)
<b>Olefins</b>		
1-nonene	720*	(1)
1-hexadecene	0.352*	(1)
<b>Aromatics</b>		
n-propylbenzene	456*	(1)
naphthalene	11.3*	(1)
n-decylbenzene	0.171*	(1)
3-hexylnaphthalene	0.072	(1)
(1) US EPA, 2000 (MPBPWIN, EPI Suite V3.10) Note: Model input data signified by "*" indicates the value was cited by EPI-Suite™ as being from the experimental database (EPA 2000).		

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

Model Input (log K<sub>ow</sub> :)

	Log Kow	Reference
<b>Paraffins (n-, iso-)</b>		
n-nonane	4.8	(1)
iso-nonane	4.7	(1)
n-hexadecane	8.2	(1)
iso-hexadecane	8.1	(1)
<b>Naphthenes</b>		
propylcyclohexane	4.6	(1)
bicyclo[4.3.0]nonane	3.7	(1)
n-decylcyclohexane	8.0	(1)
2-hexyl[4.4.0]bicyclodecane	7.1	(1)
<b>Olefins</b>		
1-nonene	5.2*	(1)
1-hexadecene	8.1	(1)
<b>Aromatics</b>		
n-propylbenzene	3.7*	(1)
naphthalene	3.3*	(1)
n-decylbenzene	7.4*	(1)
3-hexylnaphthalene	6.2	(1)

(1) US EPA, 2000 (KOWWIN, EPI Suite V3.10)  
Note: Model input data signified by "\*" indicates the value was cited by EPI-Suite™ as being from the experimental database (EPA 2000).

Model Input ( Melting Point: )

	Melting Pt °C	Reference
<b>Paraffins (n-, iso-)</b>		
n-nonane	-53.5*	(1)
iso-nonane	-80.3*	(1)
n-hexadecane	18.1*	(1)
iso-hexadecane	1.7	(1)
<b>Naphthenes</b>		
propylcyclohexane	-94.9*	(1)
bicyclo[4.3.0]nonane	-53.0*	(1)
n-decylcyclohexane	-1.7*	(1)
2-hexyl[4.4.0]bicyclodecane	23.9	(1)
<b>Olefins</b>		
1-nonene	-81.3*	(1)
1-hexadecene	4.1*	(1)
<b>Aromatics</b>		
n-propylbenzene	-99.5*	(1)
naphthalene	80.2*	(1)
n-decylbenzene	-14.4	(1)
3-hexylnaphthalene	-3*	(1)

(1) US EPA, 2000 (MPBPWIN, EPI Suite V3.10)  
Note: Model input data signified by "\*" indicates the value was cited by EPI-Suite™ as being from the experimental database (EPA 2000).

Henry's Law Constant :

Calculated by EQC for each constituent

Model Concentration -- Air :

Model Concentration --

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

Water :

Model Concentration --  
Soil :

Model Concentration --  
Sediment :

Results Remarks :

#### STUDY/METHOD

Key Study Sponsor  
Indicator : Key

Year Study Performed  
:

Method/Guideline  
Followed : EQC-Equilibrium Criterion Model, Fugacity-Based Level 1

Deviations from  
Method/Guideline :

Method/Guideline  
Description :

The EQC model calculates the distribution of a fixed quantity of conserved (i.e., non-reacting) chemical, in a closed environment at equilibrium, with no degrading reactions, no advective processes, and no intermedia transport processes (e.g., no wet deposition or sedimentation). The medium receiving the emission is unimportant because the chemical is assumed to become instantaneously distributed.

Method/Guideline and  
Test Condition  
Remarks :

GLP : No

Trent University. 2003. EQC fugacity-based EQC-equilibrium criterion model, Version 2.02. Canadian Environmental modeling Centre, Trent University, Ontario. URL:  
<http://www.trentu.ca/cemc/>

Study Reference : U.S. EPA. 2000. EPI (Estimation Programs Interface) Suite, V3.10. U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics, Washington, DC.

Yaws, C.L., X. Pan, and X. Lin. 1993. Water solubility data for 151 hydrocarbons. Chem. Eng. Feb. 100(2):108-111.

#### RELIABILITY/DATA QUALITY

Reliability : (2) Reliable with restrictions

Reliability Remarks : Environmental distribution was estimated using an accepted validated model.

#### 3.5 BIODEGRADATION

Type : Aerobic  
Inoculum : Activated sludge  
Contact time : 28 day(s)  
Method : OECD Guide-line 301 F "Ready Biodegradability: Manometric  
Respirometry Test"  
Year : 1999

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

- GLP** : No  
**Test substance** : Nigerian Diesel Fuel; CAS No. 68334-30-5
- Remark** : This diesel fuel stream did not satisfy the test criteria for ready degradability of 60% degradability within 28 days. Although this diesel fuel is not considered 'readily' biodegradable, it is inherently biodegradable since significant degradation did occur, based on EPA guidance for using ready and inherent biodegradability tests (<http://www.epa.gov/oppt/exposure/docs/half-life.htm>.)
- Result** : Average biodegradation for duplicate test flasks at 28 days:  
diesel fuel = 57.5%  
rapeseed oil = 84.4% (satisfied positive control criteria).  
The report noted that the oxygen consumption of the blank controls was below 60 mg/l as required by the test guidelines.
- Test condition** : Activated sludge inoculum for the test was collected from Medford Municipal Wastewater Treatment Plant in Medford, NJ. The plant treats predominately domestic sewage. To reduce background oxygen consumption, the activated sludge was aerated for approximately 27 hours prior to use. Sufficient supernatant was decanted to provide a 1% (v/v) inoculum for each respirometry vessel. The sewage inoculum had a microbial density of  $1E^4$  colony forming units per ml, as measured using a commercial dip-slide method, and was within the guideline criteria of  $1E^7$ - $1E^8$  CFU/l.  
Test flasks contained 990 ml of inoculated mineral salts medium, ca. 25 mg of test substance and 10 ml of sludge supernatant (inoculum). Also included in the experiment were duplicate blank flasks containing inoculated medium and duplicate reference substance flasks containing inoculated medium and 25 mg of low erucic acid rapeseed oil (LEAR). The test was run at 22 °C ( $\pm$  1°C) using a C.E.S. Aerobic Respirometer for 28 days. The instrument determined the oxygen demand every hour and replenished the oxygen through the electrolysis of copper sulfate. The extent of biodegradation over 28 days was calculated as the measured biochemical oxygen demand (BOD) expressed as a percentage of the ThOD (theoretical oxygen demand).
- Reliability** : (1) valid without restriction (58)
- Type** : Aerobic  
**Method** : ASTM D2329-68  
**Year** : 1971  
**GLP** : No  
**Test substance** : Kerosene, hydrotreated Avtur (Aviation turbine fuel)
- Method** : BOD test method using a closed-system respirometer apparatus. Referenced as ASTM D2329-68
- Remark** : Note: Test design details were minimal -no information on test systems (i.e., replicates, blank control O<sub>2</sub> data) were included in the citation, no water chemistry reported, no discussion on TOD determination for substrates included. However glucose (positive control) 5 day results were within required criteria for both fresh (70%) and saltwater media (55%) containing nutrient solution. Also oleic acid, a sparingly soluble oil, but easily degradable substrate achieved comparable degradation compared to glucose, with 67% (freshwater) and 53% degradation (seawater) after 5

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

**Result** : day incubation at 30 °C. Additionally, when testing these two rapidly degradable substrates in seawater with no nitrogen or phosphorus amendments, degradation was significantly reduced at 5 days, with only 30% of the glucose and 13% of oleic acid degraded. Based on the performance for these rapidly degradable substances, the reliability of the experimental results are acceptable.

: After 10 days, kerosene was degraded up to 46% in freshwater, while kerosene was degraded in seawater up to 53% and 2%, respectively, with and without nitrogen and phosphorus amendments.

The results of biodegradation testing of kerosene in freshwater and seawater are as follows:

<u>Substrate/Medium</u>	<u>Nutrients Added</u>	<u>BOD/TOD, %</u>	
		<u>5 days</u>	<u>10 days</u>
Kerosene			
Fresh water	+	41	46
Sea water	+	36	53
Sea water	-	2	2

**Test condition** : Fresh water amended with nutrient salts as per ASTM method plus 3 ml of effluent from a laboratory scale activated sludge plant (adapted to consumption of oil) was spiked with approximately 20 - 30 mg/l kerosene. Cell counts of sludge effluent were not reported. Seawater was sampled from the North Sea about 10 miles west of Den Helder, Netherlands. Average seawater salinity was 31.8 g/l and an average of 200-500 cells/ml bacteria was determined with the membrane filter technique. Seawater experiments were run with and without amendments of nitrogen (as ammonia) and phosphorous (as phosphate) in varying concentrations of each. Fresh water experiments were run with N and P amendments according the official method. Tests in both freshwater and seawater were conducted at 30 °C.

**Conclusion** : Experimental results indicate that the rate of degradation was highly dependent on the available nitrogen and phosphorus for a variety of substrates, but was critical in influencing the extent of kerosene degradation in sea water.

**Reliability** : (2) valid with restrictions  
Acceptable, well-documented publication which meets basic scientific principles.

(33)

**Inoculum** : Activated sludge  
**Contact time** : 28 day(s)  
**Method** : OECD Guide-line 301 F "Ready Biodegradability: Manometric Respirometry Test"

**Year** : 1999

**GLP** : No

**Test substance** : Kerosene Mid-Blend; CAS No. 8008-20-6

**Remark** : This kerosene stream did not satisfy the test criteria for ready degradability of 60% degradability within 28 days. Although this kerosene stream is not considered 'readily' biodegradable, it is inherently biodegradable since significant degradation did occur, based on EPA guidance for using ready and inherent biodegradability tests (<http://www.epa.gov/oppt/exposure/docs/half-life.htm>.)

**Result** : Average biodegradation for duplicate test flasks at 28 days:  
kerosene = 58.6%  
rapeseed oil = 84.4% (satisfied positive control criteria).

### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

- Test condition** : The report noted that the oxygen consumption of the blank controls was below 60 mg/l as required by the test guidelines.
- : Activated sludge inoculum for the test was collected from Medford Municipal Wastewater Treatment Plant in Medford, NJ. The plant treats predominately domestic sewage. To reduce background oxygen consumption, the activated sludge was aerated for approximately 27 hours prior to use. Sufficient supernatant was decanted to provide a 1% (v/v) inoculum for each respirometry vessel. The sewage inoculum had a microbial density of  $1E^{+4}$  colony forming units per ml, as measured using a commercial dip-slide method, and was within the guideline criteria of  $1E^7$ - $1E^8$  CFU/l.
- Test flasks contained 990 ml of inoculated mineral salts medium , ca. 25 mg of test substance and 10 ml of sludge supernatant (inoculum). Also included in the experiment were duplicate blank flasks containing inoculated medium and duplicate reference substance flasks containing inoculated medium and 25 mg of low erucic acid rapeseed oil (LEAR). The test was run at 22 °C ( $\pm$  °C) using a C.E.S. Aerobic Respirometer for 28 days. The instrument determined the oxygen demand every hour and replenished the oxygen through the electrolysis of copper sulfate. The extent of biodegradation over 28 days was calculated as the measured biochemical oxygen demand (BOD) expressed as a percentage of the ThOD (theoretical oxygen demand).
- Reliability** : (1) valid without restriction

(59)



### 3. Environmental Fate and Pathways

Id Kerosene  
Date 9/21/2010

#### 3.8 ADDITIONAL REMARKS

**Memo** : Biodegradability of kerosene/jet fuels

**Remark** : Removal of dissolved hydrocarbon components in water soluble fractions of aviation jet fuel was >99% for two dilution rates in a chemostat culture.

Biodegradation of water soluble fractions of kerosene-type aviation fuel (e.g., Jet A, Jet A1, and JP-8) were measured in a chemostat culture apparatus. A mixed microbial culture was isolated from groundwater contaminated with aviation fuel. Inoculum for the experiments was pre-grown for three days in culture medium of basal medium M9 diluted 10-fold by distilled water plus trace elements and fuel (1% v/v). Feeding medium for the chemostat was prepared using basal medium M9 diluted 10-fold by distilled water and autoclaved. Jet fuel was added after autoclaving in the ratio 5:1000 (0.5%). The mixture was stirred for 5 hours, and the insoluble part was separated and discarded. Cultivations were carried out in 2-l fermentators. The ratio of inoculation was 1:10, with the volume in the fermentator maintained at 1 l by an overflow device. Experiments were run for 113 hours using fermentator volumes of 11.3 and 19.4 for dilution rates of 0.1/h and 0.17/h, respectively. An abiotic control was run under the same environmental conditions except that the solution held sodium azide (2000 mg/l) to prevent bacterial growth. Experiments were run at 27 °C. The concentrations of soluble hydrocarbons before and after treatment in the chemostat were measured by GC/MS.

Residual nonpolar hydrocarbon concentrations were 2.1 µg/l and 0.42 µg/l in experiments with dilution rates of 0.17/h and 0.1/h, respectively. The measured nonpolar hydrocarbon concentration in the chemostat culture medium was 4.78 mg/l. Removal achieved >99% for both dilution rates.

**Source** : Rozkov, A., A. Kaard, and R. Vilu (1998) Biodegradation of dissolved jet fuel in chemostat by a mixed bacterial culture isolated from a heavily polluted site. Biodegradation. 8:363-369.

**Reliability** : (2) valid with restrictions

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

### 4.1 ACUTE/PROLONGED TOXICITY TO FISH

**Type** : Semistatic  
**Species** : Oncorhynchus mykiss (Fish, fresh water)  
**Exposure period** : 96 hour(s)  
**Unit** : mg/l  
**Limit test** : No  
**Analytical monitoring** : Yes  
**Method** : OECD Guide-line 203 "Fish, Acute Toxicity Test"  
**Year** : 1995  
**GLP** : Yes  
**Test substance** : CAS No. 91770-15-9; Kerosene (petroleum), sweetened

**Method** : LL<sub>50</sub> values calculated using binomial and Trimmed Spearman-Kärber methods; the NOEL was determined using Dunnett's Test.

**Remark** : The report did not specify the minimum level of detection for the analytical measurements, but the lowest calibration standard was 0.66 mg/l. Given this, the measurements at the nominal level of 4.5 mg/l and below were extrapolations outside the calibration range.

**Result** : At the end of the test, all fish of the control, and the 0.36, 0.9, and 4.5 mg/l WAF solutions were alive and showed no abnormal effects. Fish exposed to the 23 mg/l WAF solution had 53% mortality with abnormal effects of dark pigmentation and/or lethargic in the surviving fish. All fish exposed to the 50 mg/l WAF solution died during the first 24 hours of the test.

The LL<sub>50</sub> values with confidence intervals were:

<b>Exposure Time, hours</b>	<b>LL<sub>50</sub>, mg/l</b>	<b>Confidence Interval, mg/l</b>
24	36	23 - 50
48	23	17 - 30
72	19	14 - 26
96	18	13 - 24

The maximum nominal loading rate causing no mortality was 4.5 mg/l.  
The minimum nominal loading rate causing 100% mortality was 50 mg/l.  
The No-Observed-Effect Level (NOEL) was 4.5 mg/l.

Chemical analyses of the test solutions showed (mg/l as naphthalene):

	<b>Treatment level</b>					
	<b>Control</b>	<b>0.36</b>	<b>0.9</b>	<b>4.5</b>	<b>23</b>	<b>50</b>
Day 0						
new	.0099	.0078	.015	.30	1.6	--
Day 1						
old	.0084	.0073	.015	.027	1.5	3.4
new	.017	.011	.022	.43	1.9	--
Day 2						
old	ND	ND	.033	.33	1.7	--
new	ND	ND	.22	.41	1.8	--
Day 3						
old	ND	ND	.12	.36	1.6	--
new	ND	.039	.05	.41	1.9	--
Day 4						
old	ND	ND	.017	.3	1.2	--

ND = not detected

-- means all fish dead, no further measurements were made.

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

### Test condition

Deviations were noted for:

- 1) to maintain dissolved oxygen in close vessel, slightly smaller fish than guideline recommendations were used,
- 2) temperature 0.3 °C above limit for a brief time during the holding period,
- 3) temperature 0.4 °C above maximum limit for 5 hours during testing
- 4), due to a calculation error, one of the test levels was 0.36 mg/l rather than the intended 0.2 mg/l,
- 5) test substance was not held under refrigerated storage for approximately 3 weeks.

: Daily renewal of the test media ensured that test material levels were maintained at the exposure concentrations. Test solutions were prepared as water accommodated fractions (WAFs) with each treatment level prepared independently of other treatment levels. To prepare the WAFs, an appropriate volume of test substance was added quantitatively via syringe to glass carboys holding dilution water. The glass carboys were stoppered then the mixtures stirred for 24 hours ( 10% vortex, approximately 1-2 cm). An equilibration study run prior to the test justified a 24-hour stirring period. After a 1-hour settling period, WAF solutions were siphoned from the bottom of the carboy into glass test chambers. Test chambers were 4-liter (nominal) glass aspirator bottles. Bottles were completely filled with no headspace when used in the test. Three replicate test bottles were prepared for each treatment level. Nominal loading rates used in the test were 0 (control), 0.36, 0.9, 4.5, 23, and 50 mg/l. No undissolved test substance was observed in any of the WAF solutions during the test.

Test fish were juvenile rainbow trout (approximately 5 weeks old) originating from a commercial supplier (Thomas Fish Company, Anderson, CA) and fed a commercial fish diet (Ziegler Bros., Inc., Gardners, PA and Tetramin®) during the holding period. They were held in dilution water at least 12 days and at approximately 15 °C for at least 7 days prior to use in testing. Mortality during the holding period was insignificant. Food was withheld approximately 24 hours before testing. Fish used in the test measured 2.8 cm (sd = 0.3) mean total length and 0.162 g (sd = 0.05) mean weight.

To start the test, individual fish were randomly selected one at a time and placed into intermediate holding chambers, one for each test chamber, until each holding chamber contained five fish. Fish were then transferred to their respective test chambers, which then were sealed with no headspace. Biomass loading in the test chambers was 0.18 g/l.

Samples of the WAF solutions were collected on Days 0, 1, 2, 3, and 4 for chemical analysis. Freshly-prepared WAFs were collected from the mixing vessels while 24-hour old WAFs were collected as a composite from all replicates of a treatment level. WAF samples were extracted with hexane and analyzed by Gas Chromatography using a Flame Ionization Detector (GC-FID) and a capillary column. Measured concentrations were presented in the report as mg/l as naphthalene.

The test was conducted in a temperature-controlled area under a 16 hour/8 hour, light/dark photoperiod. Daylight intensity ranged from 70 to 71 foot-candles. Measurements of fresh solutions for water quality parameters showed dissolved oxygen concentration ranged from 8.3 to 9.9 mg/l, pH ranged from 7.9 to 8.7, and temperature ranged from 14 to 15 °C. Measurements of old solutions showed dissolved oxygen concentration

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

ranged from 6.8 to 8.5 mg/l and pH ranged from 7.7 to 8.3. The temperature of the old solutions was not measured.

Dilution water was a mixture of carbon filtered well water and water dialyzed by reverse osmosis. This water had an alkalinity of 63 - 65 mg CaCO<sub>3</sub>/l, hardness of 180 - 220 mg CaCO<sub>3</sub>/l, specific conductance of 380 - 410 µmhos, pH of 7.5 - 7.7, and a dissolved oxygen concentration of 8.7 - 10 mg/l at the beginning and end of the test. Total organic carbon measured 1.3 ppm during the monthly screening.

**Reliability** : (1) valid without restriction (48)

**Type** : Semistatic  
**Species** : Oncorhynchus mykiss (Fish, fresh water)  
**Exposure period** : 96 hour(s)  
**Unit** : mg/l  
**Limit test** : No  
**Analytical monitoring** : Yes  
**Method** : OECD Guide-line 203 "Fish, Acute Toxicity Test"  
**Year** : 1995  
**GLP** : Yes  
**Test substance** : CAS No. 64742-81-0; kerosene (petroleum), hydrodesulfurized

**Method** : LL<sub>50</sub> values calculated using the graphical and binomial methods; the NOEL was determined using Dunnett's Test.

**Remark** : The report did not specify the minimum level of detection for the analytical measurements, but the lowest calibration standard was 0.66 mg/l. Given this, the measurements at the nominal level of 4.5 mg/l and below were extrapolations outside the calibration range.

**Result** : At the end of the test, all fish of the control, and the 0.3, and 1.4 mg/l WAF solutions were alive and showed no abnormal effects. Fish exposed to the 6.8 mg/l WAF solution had 0% mortality with abnormal effects of dark pigmentation or erratic swimming in 27% of the surviving fish. Seventy-three percent of the fish showed no abnormal effects. Fish exposed to the 34 mg/l WAF solution had 100% mortality by 72 hours. Fish exposed to the 75 mg/l WAF solution had 93% mortality by 96 hours, with one surviving fish showing abnormal effects of dark pigmentation and lethargic.

The LL<sub>50</sub> values with confidence intervals were:

<b>Exposure Time, hours</b>	<b>LL<sub>50</sub>, mg/l</b>	<b>Confidence Interval, mg/l</b>
24	16	could not calculate
48	16	could not calculate
72	20	6.8 - 34
96	20	6.8 - 34

The maximum nominal loading rate causing no mortality was 6.8 mg/l.  
The minimum nominal loading rate causing 100% mortality was 34 mg/l.  
The No-Observed-Effect Level (NOEL) was 6.8 mg/l.

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

Chemical analyses of the test solutions showed (mg/l as naphthalene):

	Treatment level					
	Control	0.3	1.4	6.8	34	75
Day 0						
new	.0091	.047	.13	.12	4.6	3
Day 1						
old	.099	.034	.17	.66	4	2.6
new	.0091	.017	.12	.83	3.1	2.5
Day 2						
old	.014	.0023	.063	.68	2.8	2.5
new	.0021	.0029	.11	.77	1.8	5.4
Day 3						
old	.072	.072	.12	.83	1.2	5.7
new	.014	.039	.15	.53	2.8	4.6
Day 4						
old	.0025	.002	.038	.59	--	4.5

ND = not detected

-- = All fish dead at this concentration, no measurements taken.

Deviations were noted for:

- 1) to maintain dissolved oxygen in close vessel, slightly smaller fish than guideline recommendations were used
- 2) temperature excursions of 0.1 and 0.2 °C above limit for two brief times during the holding period
- 3) dissolved oxygen concentrations fell below 60% saturation on several occasion during the definitive test. The deviation ranged from 0.5 to 1.1 mg/l below the minimum limit (5.9 mg/l at 16 °C)
- 4) test substance was not held under refrigerated storage for approximately 3 weeks
- 5) biological observations were made 7 minutes outside the timeframe of  $\pm 1$  hour
- 6) the loading rates assigned for the definitive test were not in a contiguous geometric series.

### Test condition

- : Daily renewal of the test media ensured that test material levels were maintained at the exposure concentrations. Test solutions were prepared as water accommodated fractions (WAFs) with each treatment level prepared independently of other treatment levels. To prepare the WAFs, an appropriate volume of test substance was added quantitatively via syringe to glass carboys holding dilution water. The glass carboys were stoppered then the mixtures stirred for 24 hours ( 10% vortex, approximately 1-2 cm). An equilibration study run prior to the test justified a 24-hour stirring period. After a 1-hour settling period, WAF solutions were siphoned from the bottom of the carboys into glass test chambers. Test chambers were 4-liter (nominal) glass aspirator bottles. Bottles were completely filled with no headspace when used in the test. Three replicate test bottles were prepared for each treatment level. Nominal loading rates used in the test were 0 (control), 0.3, 1.4, 6.8, 34, and 75 mg/l. No undissolved test substance was observed in any of the WAF solutions during the test.

Test fish were juvenile rainbow trout (approximately 6.5 weeks old) originating from a commercial supplier (Thomas Fish Company, Anderson, CA) and fed a commercial fish diet (Ziegler Bros., Inc., Gardners, PA and Tetramin®) during the holding period. They were held in dilution water at least 12 days and at approximately 15 °C for at least 7 days prior to use in

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

testing. Mortality during the holding period was insignificant. Food was withheld approximately 24 hours before testing. Fish used in the test measured 3.6 cm (sd = 0.2) mean total length and 0.377 g (sd = 0.057) mean weight.

To start the test, individual fish were randomly selected one at a time and placed into intermediate holding chambers, one for each test chamber, until each holding chamber contained five fish. Fish were then transferred to their respective test chambers, which then were sealed with no headspace. Biomass loading in the test chambers was 0.42 g/l.

Samples of the WAF solutions were collected on Days 0, 1, 2, 3, and 4 for chemical analysis. Freshly-prepared WAFs were collected from the mixing vessels while 24-hour old WAFs were collected as a composite from all replicates of a treatment level. WAF samples were extracted with hexane and analyzed by Gas Chromatography using a Flame Ionization Detector (GC-FID) and a capillary column. Measured concentrations were presented in the report as mg/l as naphthalene.

The test was conducted in a temperature-controlled area under a 16 hour/8 hour, light/dark photoperiod. Daylight intensity ranged from 67 to 68 foot-candles. Measurements of fresh solutions for water quality parameters showed dissolved oxygen concentration ranged from 8.0 to 9.8 mg/l, pH ranged from 7.5 to 8.7, and temperature ranged from 15.6 to 16.0 °C. Measurements of old solutions showed dissolved oxygen concentration ranged from 4.8 to 8.0 mg/l and pH ranged from 7.1 to 8.1. The temperature of the old solutions was not measured.

Dilution water was a mixture of carbon filtered well water and water dialyzed by reverse osmosis. This water had an alkalinity of 64 - 70 mg CaCO<sub>3</sub>/l, hardness of 170 - 174 mg CaCO<sub>3</sub>/l, specific conductance of 380 - 420 µmhos, pH of 7.6, and a dissolved oxygen concentration of 8.5 - 8.8 mg/l at the beginning and end of the test. Total organic carbon measured 1.3 ppm during the monthly screening.

**Reliability** : (1) valid without restriction (52)

**Type** : Semistatic  
**Species** : Oncorhynchus mykiss (Fish, fresh water)  
**Exposure period** : 96 hour(s)  
**Unit** : mg/l  
**Limit test** : No  
**Analytical monitoring** : Yes  
**Method** : OECD Guide-line 203 "Fish, Acute Toxicity Test"  
**Year** : 1994  
**GLP** : Yes  
**Test substance** : CAS No. 64742-81-0, Kerosene, hydrotreated

**Method** : Visual inspection  
**Result** : 96-hr LL<sub>50</sub> lies between 10-100 mg/l WAF, based on nominal loading rates. Mortality at 96 hrs was 0, 0, 100, and 100% in the 0, 10, 100, and 1000 mg/l treatments.  
Only four concentrations were tested which is less than a minimum of five concentrations stated in the guidelines. Water hardness was higher than targeted range of 50 - 250 mg/l as CaCO<sub>3</sub>. Hardness range of 284 - 288 mg/l as CaCO<sub>3</sub> is normal for this laboratory and does not affect the survival of the fish.

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

- Analytical method used was gas chromatography-mass spectrometry. Mean reduction in the concentration of dissolved components of kerosene for each 24-hr period was 40%, 19%, and 7% for the 10, 100, and 1000 mg/l WAFs.
- Test condition** : Individual treatment concentrations were prepared as water accommodated fractions (WAF). Nominal loading rates in the definitive test were 0, 10, 100, and 1000 mg/l. Control and dilution water was laboratory mains tap water obtained from bore holes, and passed through particle and activated carbon filters (alkalinity 252 mg/l as CaCO<sub>3</sub>, hardness 277 mg/l as CaCO<sub>3</sub>, conductivity 520 S/cm, pH 7.4). Test substance was mixed in dilution water for ~70 hrs in sealed vessels with minimal headspace. Mixing time was determined in an equilibration study in which the test substance concentration in the aqueous phase of the WAFs was monitored by GC-MS. Mixtures were allowed to settle 1 hr prior to drawing off the aqueous phase for testing. Test vessels were sealed 12-liter glass aspirators completely filled with test solution and contained 7 fish per vessel. Test fish had a mean length of 4.5 cm and a mean weight of 0.83 g. Fingerlings were obtained from Zeals Trout Farm, Zeals, Wiltshire, U.K, and acclimated to test conditions for more than 9 days before use. One replicate per treatment and control were used. Test solutions were renewed daily with surviving fish transferred to the freshly prepared WAFs. Test temperature was 14 - 15 °C. Photoperiod was 16 hrs light and 8 hrs dark. Dissolved oxygen was >60% saturation (8.5 to 9.8 mg/l). pH was 7.3 - 7.8.
- To monitor the concentration of dissolved components in the test solutions, samples were collected at the beginning and end of each 24-hr period for each of the batches of WAFs prepared during the 96-hr test.
- Reliability** : (1) valid without restriction (70)
- Type** : Semistatic
- Species** : *Oncorhynchus mykiss* (Fish, fresh water)
- Exposure period** : 96 hour(s)
- Unit** : mg/l
- Limit test** : No
- Analytical monitoring** : Yes
- Method** : OECD Guide-line 203 "Fish, Acute Toxicity Test"
- Year** : 1995
- GLP** : Yes
- Test substance** : CAS No. 101316-80-7; Solvent naphtha (petroleum), hydrocracked heavy aromatic
- Method** : Statistical method:  
LL<sub>50</sub> values calculated using the probit method; the NOEL was determined using Dunnett's Test.
- Remark** : The report did not specify the minimum level of detection for the analytical measurements, but the lowest calibration standard was 0.66 mg/l. Given this, most of the measurements at the nominal level of 6.8 mg/l and below were extrapolations outside the calibration range.
- Result** : At the end of the test, all fish of the control, and the 0.3, and 1.4 mg/l WAF solutions were alive and showed no abnormal effects. Fish exposed to the 6.8 mg/l WAF solution had 6.7% mortality with abnormal effects of dark pigmentation and/or lethargic in 50% of the surviving fish. Fifty percent of the surviving fish showed no abnormal effects. Fish exposed to the 34 mg/l WAF solution had 80% mortality with abnormal effect of lethargic in all surviving fish. Fish exposed to the 75 mg/l WAF solution had 95% mortality by 48 hours, and all fish were dead by the end of the test.

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

The LL<sub>50</sub> values with confidence intervals were:

<u>Exposure Time, hours</u>	<u>LL<sub>50</sub> mg/l</u>	<u>Confidence Interval, mg/l</u>
24	could not calculate	
48	42	33 - 53
72	38	30 - 49
96	25	20 - 32

The maximum nominal loading rate causing no mortality was 1.4 mg/l.  
The minimum nominal loading rate causing 100% mortality was 75 mg/l.  
The No-Observed-Effect Level (NOEL) was 6.8 mg/l.

Chemical analyses of the test solutions showed (mg/l as naphthalene):

	<b>Treatment level</b>					
	<b>Control</b>	<b>0.3</b>	<b>1.4</b>	<b>6.8</b>	<b>34</b>	<b>75</b>
Day 0						
new	.029	.0026	.024	.5	1.9	2.8
Day 1						
old	.0014	.00073	.014	.35	1.7	2.7
new	ND	.00074	.083	.32	2.1	3.2
Day 2						
old	.0058	.069	.032	.29	1.7	2.9
new	.029	.041	.035	.91	2.4	3.2
Day 3						
old	.021	.012	.047	.61	1.4	2.9
new	.0014	.0021	.031	.47	1.9	3.2
Day 4						
old	.031	.0091	.041	.41	1.8	3.3

ND = not detected

Deviations were noted for:

- 1) on day 2, a 0.1 mg/l solution was used to renew the 0.3 mg/l treatment. This was observed and corrected within 30-45 minutes
- 2) to maintain dissolved oxygen in close vessel, slightly smaller fish than guideline recommendations were used
- 3) temperature 0.1 °C above limit for a brief time during the holding period
- 4) Dissolved oxygen concentrations fell below 60% saturation on several occasions during the definitive test. The deviation ranged from 0.1 to 0.7 mg/l below the minimum limit (5.9 mg/l at 16 °C)
- 5) test substance was not held under refrigerated storage for approximately 3 weeks
- 6) the loading rates assigned for the definitive test were not in a contiguous geometric series.

### Test condition

- : Daily renewal of the test media ensured that test material levels were maintained at the exposure concentrations. Test solutions were prepared as water accommodated fractions (WAFs) with each treatment level prepared independently of other treatment levels. To prepare the WAFs, an appropriate volume of test substance was added quantitatively via syringe to glass carboys holding dilution water. The glass carboys were stoppered then the mixtures stirred for 24 hours ( 10% vortex, approximately 1-2 cm). An equilibration study run prior to the test justified a 24-hour stirring period. After a 1-hour settling period, WAF solutions were siphoned from the bottom of the carboys into glass test chambers. Test chambers were 4-liter (nominal) glass aspirator bottles. Bottles were completely filled with no



headspace when used in the test. Three replicate test bottles were prepared for each treatment level. Nominal loading rates used in the test were 0 (control), 0.3, 1.4, 6.8, 34, and 75 mg/l. No undissolved test substance was observed in any of the WAF solutions during the test.

Test fish were juvenile rainbow trout (approximately 6.5 weeks old) originating from a commercial supplier (Thomas Fish Company, Anderson, CA) and fed a commercial fish diet (Ziegler Bros., Inc., Gardners, PA and Tetramin®) during the holding period. They were held in dilution water at least 12 days and at approximately 15 C for at least 7 days prior to use in testing. Mortality during the holding period was insignificant. Food was withheld approximately 24 hours before testing. Fish used in the test measured 3.9 cm (sd = 0.4) mean total length and 0.432 g (sd = 0.113) mean weight.

To start the test, individual fish were randomly selected one at a time and placed into intermediate holding chambers, one for each test chamber, until each holding chamber contained five fish. Fish were then transferred to their respective test chambers, which then were sealed with no headspace. Biomass loading in the test chambers was 0.48 g/l.

Samples of the WAF solutions were collected on Days 0, 1, 2, 3, and 4 for chemical analysis. Freshly-prepared WAFs were collected from the mixing vessels while 24-hour old WAFs were collected as a composite from all replicates of a treatment level. WAF samples were extracted with hexane and analyzed by Gas Chromatography using a Flame Ionization Detector (GC-FID) and a capillary column. Measured concentrations were presented in the report as mg/l as naphthalene.

The test was conducted in a temperature-controlled area under a 16 hour/8 hour, light/dark photoperiod. Daylight intensity ranged from 55 to 68 foot-candles. Measurements of fresh solutions for water quality parameters showed dissolved oxygen concentration ranged from 7.8 to 9.6 mg/l, pH ranged from 7.8 to 8.7, and temperature measured a constant 16 °C. Measurements of old solutions showed dissolved oxygen concentration ranged from 5.2 to 7.8 mg/l and pH ranged from 7.1 to 8.2. The temperature of the old solutions was not measured.

Dilution water was a mixture of carbon filtered well water and water dialyzed by reverse osmosis. This water had an alkalinity of 64 - 70 mg CaCO<sub>3</sub>/l, hardness of 170 - 174 mg CaCO<sub>3</sub>/l, specific conductance of 380 - 420 µmhos, pH of 7.6, and a dissolved oxygen concentration of 8.5 - 8.8 mg/l at the beginning and end of the test. Total organic carbon measured 1.3 ppm during the monthly screening.

**Reliability** : (1) valid without restriction

(50)

#### 4.2 ACUTE/PROLONGED TOXICITY TO AQUATIC INVERTEBRATES

**Type** : Static renewal  
**Species** : Daphnia magna (Crustacea)  
**Exposure period** : 48 hour(s)  
**Unit** : mg/l  
**Limit Test** : no  
**Analytical monitoring** : yes  
**Method** : OECD Guide-line 202

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

- Year** : 1995  
**GLP** : yes  
**Test substance** : CAS No. 91770-15-9; Kerosene (petroleum), sweetened
- Method** : EL<sub>50</sub> values calculated using graphical and probit methods, the NOEL was determined by Dunnett's Test.
- Remark** : The report did not specify the minimum level of detection for the analytical measurements, but the lowest calibration standard was 0.66 mg/l. Given this, the measurements at the nominal level of 4.5 mg/l and below were extrapolations outside the calibration range.
- Result** : At the end of the test, one individual from the control group was immobile, all other daphnids of the control group appeared normal. Daphnids exposed to the 0.1, 0.36, and 0.9 mg/l WAF solutions were alive and appeared normal. Daphnids exposed to the 4.5 mg/l WAF solution had 15% immobilization, with surviving daphnids appearing normal. Daphnids exposed to the 23 mg/l WAF solution had 55% immobilization with surviving daphnids appearing normal. All daphnids exposed to the 50 mg/l WAF solution were immobile at the end of the test.

The EL<sub>50</sub> values with confidence intervals were:

<b>Exposure Time, hours</b>	<b>EL<sub>50</sub>, mg/l</b>	<b>Confidence Interval, mg/l</b>
24	46	could not be calculated
48	21	17 - 27

The maximum nominal loading causing no immobilization was 0.9 mg/l. The minimum nominal loading rate causing 100% immobilization was 50 mg/l. The No-Observed-Effect Level (NOEL) was 4.5 mg/l based on immobilization.

Chemical analyses of the test solutions showed (mg/l as naphthalene):

<b>Treatment Level</b>	<b>Day 0 New</b>	<b>Day 2 Old</b>
Control	ND	0.0057
0.1	ND	ND
0.36	0.0390	ND
0.9	0.050	ND
4.5	0.41	0.13
23	1.9	0.083
50	3.2	1.7

ND = not detected

Deviations were noted for:

- 1) test substance was not held under refrigerated storage for approximately 3 weeks
- 2) not all test chambers contained five daphnids; four replicates had six and one replicate had four
- 3) due to a calculation error, a loading rate of 0.36 mg/l was used instead of the intended 0.2 mg/l
- 4) test daphnids were added directly to the test vessels rather than into intermediate vessels prior to adding to the test vessels
- 5) the dilution water was not aerated prior to use.

- Test condition** : Daily renewal of the test media ensured that test material levels were maintained at the exposure concentrations. Test solutions were prepared as water accommodated fractions (WAFs) with each treatment level prepared independently of other treatment levels. To prepare the WAFs,

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

an appropriate volume of test substance was added quantitatively via syringe to glass carboys holding dilution water. The glass carboys were stoppered then the mixtures stirred for 24 hours ( 10% vortex, approximately 1-2 cm). An equilibration study run prior to the test justified a 24-hour stirring period. After a 1-hour settling period, WAF solutions were siphoned from the bottom of the carboy into glass test vessels. Test vessels were 125-ml glass Erlenmeyer flasks that were completely filled (no headspace) with the WAF solutions. Each test vessel was stoppered with glass closures to minimize evaporation and/or volatilization. Four replicate test vessels were prepared for each treatment level. Nominal loading rates used in the test were 0 (control), 0.1, 0.36, 0.9, 4.5, 23, and 50 mg/l. No undissolved test substance was observed in any of the WAF solutions during the test.

Daphnid cultures were maintained at the testing laboratory in dilution water supplemented with vitamin B12 and selenium. They were fed daily a green alga (*Selenastrum capricornutum*) and a yeast/salmon starter/cereal leaves mixture. Daphnids used in the test were 24 hours old that were taken from 17-day old parents.

To start the test, individual daphnids were randomly selected and distributed one at a time to each test vessel until each replicate test vessel contained five daphnids. Organism loading was approximately 1 daphnid per 28 ml of solution.

Samples of the WAF solutions were collected on Days 0 and 2 for chemical analysis. Day 0 samples were the freshly-prepared WAFs, while the Day 2 samples were collected as a composite from all replicates of a treatment level. WAF samples were extracted with hexane and analyzed by Gas Chromatography using a Flame Ionization Detector (GC-FID) and a capillary column. Measured concentrations were presented in the report as mg/l as naphthalene.

The test was conducted in a temperature-controlled waterbath under a 16 hour/8 hour light/dark photoperiod. Daylight intensity ranged from 68 to 70 foot-candles. Measurements of fresh solutions (Day 0) for water quality parameters showed dissolved oxygen concentration ranged from 8.3 to 9.4 mg/l, pH ranged from 7.5 to 8.4 and temperature was a constant 21 °C. Measurements of old test solutions (Day 2) indicated dissolved oxygen concentration ranged from 7.0 to 8.5, pH ranged from 7.7 to 8.2, and temperature was a constant 20 °C.

Dilution water was a mixture of carbon filtered well water and water dialyzed by reverse osmosis. This water had an alkalinity of 63 - 70 mg CaCO<sub>3</sub>/l, hardness of 170 - 180 mg CaCO<sub>3</sub>/l, specific conductance of 380 µmhos, pH of 7.6 - 7.7, and a dissolved oxygen concentration of 8.8 - 10.2 mg/l during the testing period. Total organic carbon measured 1.3 ppm during the monthly screening.

**Reliability** : (1) valid without restriction (44)

**Type** : Static renewal

**Species** : *Daphnia magna* (Crustacea)

**Exposure period** : 48 hour(s)

**Unit** : mg/l

**Limit Test** : No

**Analytical monitoring** : Yes

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

**Method** : OECD Guide-line 202  
**Year** : 1995  
**GLP** : Yes  
**Test substance** : CAS No. 64741-81-0; kerosene (petroleum), hydrodesulfurized

**Method** : EL<sub>50</sub> values calculated using the Trimmed Spearman-Karber and probit methods, NOEL determined using Dunnett's Test.

**Remark** : The report did not specify the minimum level of detection for the analytical measurements, but the lowest calibration standard was 0.66 mg/l. Given this, measurements (most) at the nominal level of 6.8 mg/l and below were extrapolations outside the calibration range.

**Result** : At the end of the test, all daphnids in the control group were alive and normal.

Daphnids exposed to the 0.1 mg/l WAF solution had one (5%) immobile daphnid at the end of the test with the remaining daphnids normal. Daphnids exposed to the 0.3 mg/l WAF solution had 15% immobilization, while daphnids exposed to the 1.4 mg/l WAF solution had 50% immobilization. Surviving daphnids of those two groups appeared normal. All daphnids exposed to 6.8 and 34 mg/l WAF solutions were immobile by Day 2 and Day 1, respectively, of the test.

The EL<sub>50</sub> values with confidence intervals were:

<u>Exposure</u> <u>Time, hours</u>	<u>EL<sub>50</sub>,</u> <u>mg/l</u>	<u>Confidence Interval,</u> <u>mg/l</u>
24	4.6	3.4 - 6.3
48	1.4	1.0 - 2.0

The maximum nominal loading causing no immobilization was not determined because all test levels had at least one immobile daphnid. The minimum nominal loading rate causing 100% immobilization was 6.8 mg/l. The No-Observed-Effect Level (NOEL) was 0.3 mg/l based on immobilization.

Chemical analyses of the test solutions showed (mg/l as naphthalene):

<u>Treatment</u> <u>Level</u>	<u>Day 0</u> <u>New</u>	<u>Day 2</u> <u>Old</u>
Control	0.0091	0.0065
0.1	0.0040	0.0008
0.3	0.047	0.0031
1.4	0.13	0.094
6.8	0.121	0.79
34	4.6	3.52

- 1 Report suggests material may have been lost during sampling.
- 2 Sample taken on Day 1 due to complete mortality.

Deviations were noted for:

- 1) test substance was not held under refrigerated storage for approximately 3 weeks
- 2) the test temperature deviated 0.12 °C for approximately seven minutes during the test
- 3) temperature measurements of the old solutions were not taken on Day 2
- 4) the dilution water was not aerated prior to the test
- 5) the definitive test loading rates did not follow a contiguous geometric series.

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

**Test condition** : Daily renewal of the test media ensured that test material levels were maintained at the exposure concentrations. Test solutions were prepared as water accommodated fractions (WAFs) with each treatment level prepared independently of other treatment levels. To prepare the WAFs, an appropriate volume of test substance was added quantitatively via syringe to glass carboys holding dilution water. The glass carboys were stoppered then the mixtures stirred for 24 hours ( 10% vortex, approximately 1-2 cm). An equilibration study run prior to the test justified a 24-hour stirring period. After a 1-hour settling period, WAF solutions were siphoned from the bottom of the carboy into glass test vessels. Test vessels were 125-ml glass Erlenmeyer flasks that were completely filled (no headspace) with the WAF solutions. Each test vessel was stoppered with glass closures to minimize evaporation and/or volatilization. Four replicate test vessels were prepared for each treatment level. Nominal loading rates used in the test were 0 (control), 0.1, 0.3, 1.4, 6.8, and 34 mg/l. No undissolved test substance was observed in any of the WAF solutions during the test.

Daphnid cultures were maintained at the testing laboratory in dilution water supplemented with vitamin B12 and selenium. They were fed daily a green alga (*Selenastrum capricornutum*) and a yeast/salmon starter/cereal leaves mixture. Daphnids used in the test were 24 hours old that were taken from 13 and 15-day old parents.

To start the test, individual daphnids were randomly selected and distributed one at a time to intermediate containers until each container held five daphnids. Daphnids were then transferred to their respective test vessel. Organism loading was approximately 1 daphnid per 28 ml of solution.

Samples of the WAF solutions were collected on Days 0 and 2 for chemical analysis. Day 0 samples were the freshly-prepared WAFs, while the Day 2 samples were collected as a composite from all replicates of a treatment level. WAF samples were extracted with hexane and analyzed by Gas Chromatography using a Flame Ionization Detector (GC-FID) and a capillary column. Measured concentrations were presented in the report as mg/l as naphthalene.

The test was conducted in a temperature-controlled waterbath under a 16 hour/8 hour light/dark photoperiod. Daylight intensity ranged from 67 to 68 foot-candles. Measurements of fresh solutions (Day 0) for water quality parameters showed dissolved oxygen concentration ranged from 8.0 to 9.5 mg/l, pH ranged from 8.0 to 8.7 and temperature was a constant 20 °C. Measurements of old test solutions (Day 2) indicated dissolved oxygen concentration ranged from 7.2 to 8.1 and pH ranged from 7.6 to 8.6. The temperature of the old solutions was not measured.

Dilution water was a mixture of carbon filtered well water and water dialyzed by reverse osmosis. This water had an alkalinity of 64 - 70 mg CaCO<sub>3</sub>/l, hardness of 170 - 174 mg CaCO<sub>3</sub>/l, specific conductance of 380 - 420 µmhos, pH of 7.6, and a dissolved oxygen concentration of 8.5 - 8.8 mg/l during the testing period. Total organic carbon measured 1.3 ppm during the monthly screening.

**Reliability** : (1) valid without restriction

(45)

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

**Type** : Static  
**Species** : Daphnia magna (Crustacea)  
**Exposure period** : 48 hour(s)  
**Unit** : mg/l  
**Limit Test** : No  
**Analytical monitoring** : Yes  
**Method** : OECD Guide-line 202  
**Year** : 1993  
**GLP** : Yes  
**Test substance** : CAS No. 64742-81-0; Kerosene, hydrotreated

**Method** : Visual inspection  
**Result** : 48-hr EL<sub>50</sub> lies between 40 and 89 mg/l WAF, based on nominal loading rates.

Numbers of immobilized daphnids after 48 hrs were 0, 0, 0, 0, 14, and 20 in the 0, 8.3, 18, 40, 89, and 200 mg/l treatments.

No excursions from protocol were noted.

Analytical method used was gas chromatography-mass spectrometry. Mean reduction in the concentration of dissolved components of kerosene during the test was 31%, 22%, and 34% in the 8.3, 40, and 200 mg/l WAFs. The sample collected at 0 hr for the 89 mg/l WAF was lost during analysis, thus, the percent reduction could not be determined for that treatment.

**Test condition** : Individual treatment concentrations were prepared as water accommodated fractions (WAF). Nominal loading rates in the definitive test were 0, 8.3, 18, 40, 89, and 200 mg/l. Control and dilution water was reconstituted hard water prepared by adding salts to reverse osmosis filtered water following EPA guidelines (hardness 180 mg/l as CaCO<sub>3</sub>). Test substance was mixed in dilution water for 23 hrs in sealed vessels with minimal headspace. Mixing time was determined in an equilibration study in which the test substance concentration in the aqueous phase of the WAFs was monitored by GC-MS. Mixtures were allowed to settle 1 hr prior to drawing off the aqueous phase for testing. Test vessels were sealed 150-ml Erlenmeyer flasks completely filled with test solution and contained 10 daphnids per vessel. Test daphnids were <24 hrs old and collected from cultures supplied by the testing laboratory that have been aged between 14 and 28 days. Two replicates per treatment and control were used.

Test temperature was 18 - 19 °C. Photoperiod was 16 hrs light and 8 hrs dark. Dissolved oxygen was >60% saturation (9.2 to 9.7 mg/l). pH was 7.6 - 8.2.

Samples were collected at the beginning and end of the test to monitor the concentration of dissolved components of kerosene in the test solutions.

**Reliability** : (1) valid without restriction

(69)

**Type** : Static renewal  
**Species** : Daphnia magna (Crustacea)  
**Exposure period** : 48 hour(s)  
**Unit** : mg/l  
**Limit Test** : No  
**Analytical monitoring** : Yes  
**Method** : OECD Guide-line 202  
**Year** : 1995  
**GLP** : Yes

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

- Test substance** : CAS No. 101316-80-7; solvent naphtha (petroleum), hydrocracked heavy aromatic
- Method** : Statistical method:  
EL<sub>50</sub> values calculated using the probit method, NOEL determined using Duncan's Multiple Range Test.
- Remark** : The report did not specify the minimum level of detection for the analytical measurements, but the lowest calibration standard was 0.66 mg/l. Given this, the measurements at the nominal level of 1.4 mg/l and below were extrapolations outside the calibration range.
- Result** : At the end of the test, two individuals (10%) from the control group were immobile, all other daphnids of the control group appeared normal. Daphnids exposed to the 0.1 mg/l WAF solution had one (5%) immobile daphnid at the end of the test, while all daphnids exposed to the 0.3 mg/l WAF solution were normal. Exposure to the 1.4 mg/l WAF solution caused 35% immobility with the surviving daphnids all normal. By the end of the test, all daphnids exposed to 6.8 and 34 mg/l WAF solutions were immobile.

The EL<sub>50</sub> values with confidence intervals were:

<b>Exposure Time, hours</b>	<b>EL<sub>50</sub>, mg/l</b>	<b>Confidence Interval, mg/l</b>
24	28	22 - 35
48	1.9	1.3 - 4.3

The maximum nominal loading causing no immobilization was 0.3 mg/l. The minimum nominal loading rate causing 100% immobilization was 6.8 mg/l. The No-Observed-Effect Level (NOEL) was 0.3 mg/l based on immobilization.

Chemical analyses of the test solutions showed (mg/l as naphthalene):

<b>Treatment Level</b>	<b>Day 0 New</b>	<b>Day 2 Old</b>
Control	0.029	0.0018
0.1	0.141	0.151
0.3	0.041	0.050
1.4	0.035	0.030
6.8	0.91	0.94
34	2.4	2.4

1 Report suggests high values due to contamination.

Deviations were noted for:

- 1) On Day 1 of the test biological observations were made outside the  $\pm 1$ -hour limit
- 2) temperature measurements of the old solutions were not taken on Day 2
- 3) the test temperature deviated by 0.1 °C beyond the  $\pm 1$  °C limit for the test for approximately 34 hours
- 4) test substance was not held under refrigerated storage for approximately 3 weeks
- 5) the definitive test loading rates did not follow a contiguous geometric series.

- Test condition** : Daily renewal of the test media ensured that test material levels were maintained at the exposure concentrations. Test solutions were prepared as water accommodated fractions (WAFs) with each treatment level prepared independently of other treatment levels. To prepare the WAFs,

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

an appropriate volume of test substance was added quantitatively via syringe to glass carboys holding dilution water. The glass carboys were stoppered then the mixtures stirred for 24 hours ( 10% vortex, approximately 1-2 cm). An equilibration study run prior to the test justified a 24-hour stirring period. After a 1-hour settling period, WAF solutions were siphoned from the bottom of the carboy into glass test vessels. Test vessels were 125-ml glass Erlenmeyer flasks that were completely filled (no headspace) with the WAF solutions. Each test vessel was stoppered with glass closures to minimize evaporation and/or volatilization. Four replicate test vessels were prepared for each treatment level. Nominal loading rates used in the test were 0 (control), 0.1, 0.3, 1.4, 6.8, and 34 mg/l. No undissolved test substance was observed in any of the WAF solutions during the test.

Daphnid cultures were maintained at the testing laboratory in dilution water supplemented with vitamin B12 and selenium. They were fed daily a green alga (*Selenastrum capricornutum*) and a yeast/salmon starter/cereal leaves mixture. Daphnids used in the test were 24 hours old that were taken from 13-day old parents.

To start the test, individual daphnids were randomly selected and distributed one at a time to intermediate containers until each container held five daphnids. Daphnids were then transferred to their respective test vessel. Organism loading was approximately 1 daphnid per 28 ml of solution.

Samples of the WAF solutions were collected on Days 0 and 2 for chemical analysis. Day 0 samples were the freshly-prepared WAFs, while the Day 2 samples were collected as a composite from all replicates of a treatment level. WAF samples were extracted with hexane and analyzed by Gas Chromatography using a Flame Ionization Detector (GC-FID) and a capillary column. Measured concentrations were presented in the report as mg/l as naphthalene.

The test was conducted in a temperature-controlled waterbath under a 16 hour/8 hour light/dark photoperiod. Daylight intensity ranged from 66 to 68 foot-candles. Measurements of fresh solutions (Day 0) for water quality parameters showed dissolved oxygen concentration ranged from 8.4 to 9.5 mg/l, pH ranged from 7.8 to 8.4 and temperature was a constant 20 °C. Measurements of old test solutions (Day 2) indicated dissolved oxygen concentration ranged from 7.8 to 8.2 and pH ranged from 7.6 to 8.4. The temperature of the old solutions was not measured.

Dilution water was a mixture of carbon filtered well water and water dialyzed by reverse osmosis. This water had an alkalinity of 64 - 70 mg CaCO<sub>3</sub>/l, hardness of 170 - 174 mg CaCO<sub>3</sub>/l, specific conductance of 380 - 420 µmhos, pH of 7.6, and a dissolved oxygen concentration of 8.5 - 8.8 mg/l during the testing period. Total organic carbon measured 1.3 ppm during the monthly screening.

### Reliability

: (1) valid without restriction

(46)



## 4. Ecotoxicity

Id Kerosene  
Date 9/21/2010

### 4.3 TOXICITY TO AQUATIC PLANTS E.G. ALGAE

**Species** : Selenastrum capricornutum (Algae)  
**Endpoint** : Growth rate  
**Exposure period** : 96 hour(s)  
**Unit** : mg/l  
**Limit test** : No  
**Analytical monitoring** : Yes  
**Method** : OECD Guide-line 201 "Algae, Growth Inhibition Test"  
**Year** : 1995  
**GLP** : Yes  
**Test substance** : CAS No. 64742-81-0; kerosene (petroleum), hydrodeulfurized

**Method** : EL<sub>50</sub> values calculated by the inverse interpolation method of Snedecor and Cochran (1989); NOEL value determined using Analysis of Variance.

**Remark** : The report did not specify the minimum level of detection for the analytical measurements, but the lowest calibration standard was 0.66 mg/l. Given this, the measurements at the nominal level of 4.0 mg/l and below were extrapolations outside the calibration range.

**Result** : The percent inhibition values for average specific growth rate (ASGR) and area under the growth curve (AUGC) at 72 and 96 hours for each WAF loading level were:

Loading Rate mg/l	% Inhibition Relative to Controls			
	ASGR		AUGC	
	72 H	96 H	72 H	96 H
0.4	-4.8	0	-41	-20
4.0	6.8	7.5	13	21
20	45	48	75	87
45	96	100	98	99
100	100	100	99	99

Note: negative values indicate a stimulatory effect.

The EL<sub>50</sub> values with confidence intervals were:

	EL <sub>50</sub> , mg/l	95% Confidence Limits
<b>ASGR</b>		
0 - 72 h	8.3	could not calculate
0 - 96 h	6.2	could not calculate
<b>AUGC</b>		
0 - 72 h	15	0 - 52
0 - 96 h	11	0 - 42

The No-Observed-Effect Levels determined for ASGR and AUGC were:

ASGR NOEL	72 h	=	4.0 mg/l
	96 h	=	0.4 mg/l
AUGC NOEL	72 h	=	4.0 mg/l
	96 h	=	0.4 mg/l

Chemical analyses of the test solutions showed:

Treatment Level	Measured Concentration (mg/l as naphthalene)	
	Day 0	Day 4
Control	0.015	Not detected
0.4 mg/l	0.0083	0.020
4.0	0.23	0.15
20	0.86	0.89
45	3.9	2.7
100	5.4	4.6

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

### Test condition

Deviations were noted for:

- 1) test substance was not held under refrigerated storage for approximately 3 weeks
- 2) loading rates assigned for the test were not in a contiguous geometric series.

: Test solutions were prepared as water accommodated fractions (WAFs) with each treatment level prepared independently of other treatment levels. To prepare the WAFs, an appropriate volume of test substance was added quantitatively to 1.8 liters algal nutrient media in a 2-liter glass aspirator bottle. The bottles were stoppered then stirred ( 10% vortex, approximately 1 cm) for approximately 24 hours (an equilibration study conducted prior to the test justified a 24-hour stirring period). After a 1-hour settling period, the WAF solutions were removed through an outlet at the bottom of the bottle. A portion of each WAF solution was used to condition the replicate test vessels by placing 25 ml of solution per flask and allowing the solutions to sit for approximately 5 minutes. The flasks then were swirled and the solution discarded. Each replicate test vessel was filled with 140 ml of WAF, inoculated with algae and immediately closed with a ground glass stopper. Test vessels were 125-ml glass Erlenmeyer flasks, and six replicate vessels were used in the control group, while three replicate vessels were used for each WAF solution. Nominal loading rates used in the test were 0 (control), 0.4, 4.0, 20, 45, and 100 mg/l. No undissolved test substance was observed in the test vessels during the test.

Nutrient medium was prepared according to the formulation described by Miller et al. 1978 (EPA-600/9-78-018) with additional sodium bicarbonate\* (NaHCO<sub>3</sub> at 100 mg/l) added as a source of carbon in the sealed test chambers. The nutrient medium was prepared with distilled water and reagent grade chemicals at the following concentrations:

Compound	Concentration mg/l	Compound	Concentration µg/l
NaNO <sub>3</sub>	25.5	H <sub>3</sub> BO <sub>3</sub>	185.52
MgCl <sub>2</sub> .6H <sub>2</sub> O	12.164	MnCl <sub>2</sub> .4H <sub>2</sub> O	415.38
CaCl <sub>2</sub> .2H <sub>2</sub> O	4.41	ZnCl <sub>2</sub>	3.27
MgSO <sub>4</sub> .7H <sub>2</sub> O	14.7	CoCl <sub>2</sub> .6H <sub>2</sub> O	1.428
K <sub>2</sub> HPO <sub>4</sub>	1.044	CuCl <sub>2</sub> .2H <sub>2</sub> O	0.012
NaHCO <sub>3</sub>	*see above	Na <sub>2</sub> MoO <sub>4</sub> .2H <sub>2</sub> O	7.26
		FeCl <sub>3</sub> .6H <sub>2</sub> O	159.76
		Na <sub>2</sub> EDTA.2H <sub>2</sub> O	300.0

*S. capricornutum* cultures were maintained by the testing laboratory at a temperature of 24 ± 2 °C under constant illumination of 4300 10% lux provided by cool-white fluorescent bulbs. The source of the culture was the Department of Botany, University of Texas (initial strain #1648). The algal inoculum used to start the test originated from 5-day old stock cultures in log phase growth.

To start the test, the different WAF and control solutions were placed in their respective number of replicate flasks and inoculated with *S. capricornutum* to an initial cell density of 1.0 x 10<sup>3</sup> cells/ml. The flasks were sealed and randomly positioned on a shaker table. All flasks were incubated for four days under constant illumination (4300 to 4400 Lux), continuous shaking at 125 rpm, and at a mean temperature of 23.6 °C.

Cell density was determined for each replicate flask at 24, 48, 72, and 96

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

hours. This was done by taking 3.5 ml of solution from each flasks and measuring fluorescence on a Turner Filter fluorometer and converting fluorescence readings to algal cell densities using a regression formula developed through cell counts. In order to maintain no headspace in the test flasks, the 3.5 ml aliquot of solution removed for analyses was replaced with 3.5 ml of spare WAF. The spare WAF samples were stored in closed vessels, and a new vessel was opened for each replacement interval. The replacement of 3.5 ml of WAF diluted the cell densities by approximately 2.5% on Days 2-4. The overall affect of this was considered not significant.

The pH of the control and test solutions were taken at 0 and 96 hours. The pH of the solutions ranged from 7.4 to 7.5 at the beginning of the test and 7.6 to 9.8 at the end of the test.

The EL<sub>50</sub> values were determined on the percent inhibition relative to the control values for average specific growth rate and area under the growth curve. The specific growth rates for each treatment were determined by calculating the slope of the regression line of ln(cell density) versus time. The area under the growth curves was calculated in accordance with the equations in OECD Guideline No. 201

**Reliability** : (1) valid without restriction (51)

**Species** : Selenastrum capricornutum (Algae)  
**Endpoint** :  
**Exposure period** : 72 hour(s)  
**Unit** : mg/l  
**Limit test** : no  
**Analytical monitoring** : yes  
**Method** : OECD Guide-line 201 "Algae, Growth Inhibition Test"  
**Year** : 1993  
**GLP** : Yes  
**Test substance** : CAS No. 64742-81-0, Kerosene, hydrotreated

**Method** : EL<sub>50</sub> values determined by visual inspection. Williams test used to determine NOELs.

**Result** :  
Based on nominal loading rates: 72-hr EL<sub>50</sub> (biomass) lies between 10 and 30 mg/l  
72-hr EL<sub>50</sub> (growth rate) lies between 10 and 30 mg/l  
72-hr NOEL (biomass) = 1 mg/l  
72-hr NOEL (growth rate) = 10mg/l.

<b>Nominal Conc. (mg/l)</b>	<b>72 h % Inhibition</b>	<b>72 h Mean Cell Conc. (million cells/ml)</b>
Control	n/a	0.15
0.3	6.7	0.14
1.0	20	0.12
3.0	20	0.12
10	33	0.10
30	91	0.013

n/a - Not applicable

The pH increased by more than one unit during the test as a result of good culture growth and could not be avoided. This deviation was not sufficient to invalidate the study.

Analytical method used was gas chromatography-mass spectrometry. Mean reduction in the concentration of dissolved components of kerosene

## 4. Ecotoxicity

Id Kerosene  
Date 9/21/2010

**Test condition** : during the test was 50%, 0%, 50%, and 19% in the 1, 3, 10, and 30 mg/l WAFs.  
: Individual treatment concentrations were prepared as water accommodated fractions (WAF). Nominal loading rates in the definitive test were 0, 0.3, 1, 3, 10 and 30 mg/l. Control and dilution water was algal nutrient medium prepared according to EPA guidelines except that boric acid was present at 105 g/l and sodium bicarbonate at 50 mg/l. Test substance was mixed with dilution water for 20 hrs, and the mixture was allowed to settle for 1 hr prior to drawing off the aqueous phase for testing. Test vessels were sealed 300 ml Erlenmeyer flasks completely filled with test solution. There were four flasks for each treatment and seven control flasks. Three of the four treatment and six of the seven control flasks were inoculated with algal cells to yield an initial concentration of 5000 cells/ml. Algal cells were obtained from laboratory cultures that were originally derived from a strain from American Type Culture Collection (ATCC 22662). Uninoculated flasks were used to determine particle counts without algal cells using a Coulter Multisizer. Flasks were incubated in a cooled orbital (100 cycles/min) incubator. Biomass was calculated as area under the growth curve. Test temperature was 24 - 25 °C. Lighting was continuous at ~3000 lux. The pH ranged from 7.3 - 7.5 at test initiation and 8.6 - 9.4 at test termination. Samples were collected at the beginning and end of the test to monitor the concentration of dissolved components of kerosene in the test solutions.

**Reliability** : (1) valid without restriction

(71)

**Species** : Selenastrum capricornutum (Algae)

**Exposure period** : 96 hour(s)

**Unit** : mg/l

**Limit test** : No

**Analytical monitoring** : Yes

**Method** : OECD Guide-line 201 "Algae, Growth Inhibition Test"

**Year** : 1995

**GLP** : yes

**Test substance** : CAS No. 101316-80-7; solvent naphtha (petroleum), hydrocracked heavy aromatic

**Method** : Statistical method  
EL<sub>50</sub> values calculated by the inverse interpolation method of Snedecor and Cochran (1989); NOEL value determined using Duncan's Multiple Range Test.

**Remark** : The report did not specify the minimum level of detection for the analytical measurements, but the lowest calibration standard was 0.66 mg/l. Given this, the measurements at the nominal level of 50 mg/l and below were extrapolations outside the calibration range.

**Result** : The percent inhibition values for average specific growth rate (ASGR) and area under the growth curve (AUGC) at 72 and 96 hours for each WAF loading level were:

Loading Rate mg/l	% Inhibition Relative to Controls			
	ASGR		AUGC	
	72 H	96 H	72 H	96 H
0.2	1.4	1.4	8.0	10
0.8	-3.2	-1.9	-13	-9.4
6.2	-4.4	-1.5	-34	-25
12	58	68	84	94
50	92	97	98	100

Note: negative values indicate a stimulatory effect.

## 4. Ecotoxicity

Id Kerosene  
Date 9/21/2010

The EL<sub>50</sub> values with confidence intervals were:

	<u>EL<sub>50</sub>, mg/l</u>	<u>95% Confidence Limits</u>
<b>ASGR</b>		
0 - 72 h	6.7	0.4 - 896
0 - 96 h	5.0	0.5 - 131
<b>AUGC</b>		
0 - 72 h	12	could not calculate
0 - 96 h	5.9	could not calculate

The No-Observed-Effect Levels determined for ASGR and AUGC were:

ASGR NOEL 72 and 96 hours = 6.2 mg/l  
AUGC NOEL 72 and 96 hours = 12 mg/l

Chemical analyses of the test solutions showed:

<u>Treatment</u>	<u>Measured Concentration</u>	
	<u>(mg/l as naphthalene)</u>	
Level	Day 0	Day 4
Control	0.015	ND
0.2 mg/l	ND	ND
0.8	ND	0.021
6.2	0.0059	ND
12	0.51	0.44
50	2.0	1.8

Note: the authors state that the 6.2 mg/l test level may have been dosed at 3.1 mg/l, but this could not be concluded.

Deviations were noted for:

- 1) test substance was not held under refrigerated storage for approximately 3 weeks
- 2) loading rates assigned for the test were not in a contiguous geometric series.

### Test condition

- : Test solutions were prepared as water accommodated fractions (WAFs) with each treatment level prepared independently of other treatment levels. To prepare the WAFs, an appropriate volume of test substance was added quantitatively to 1.8 liters algal nutrient media in a 2-liter glass aspirator bottle. The bottles were stoppered then stirred ( 10% vortex, approximately 1 cm) for approximately 24 hours (an equilibration study conducted prior to the test justified a 24-hour stirring period). After a 1-hour settling period, the WAF solutions were removed through an outlet at the bottom of the bottle. A portion of each WAF solution was used to condition the replicate test vessels by placing 25 ml of solution per flask and allowing the solutions to sit for approximately 5 minutes. The flasks then were swirled and the solution discarded. Each replicate test vessel was filled with 140 ml of WAF, inoculated with algae and immediately closed with a ground glass stopper. Test vessels were 125-ml glass Erlenmeyer flasks, and six replicate vessels were used in the control group, while three replicate vessels were used for each WAF solution. Nominal loading rates used in the test were 0 (control), 0.2, 0.8, 6.2, 12, and 50 mg/l. No undissolved test substance was observed in the test vessels during the test.

Nutrient medium was prepared according to the formulation described by Miller et al. 1978 (EPA-600/9-78-018) with additional sodium bicarbonate\* (NaHCO<sub>3</sub> at 100 mg/l) added as a source of carbon in the sealed test chambers. The nutrient medium was prepared with distilled water and reagent grade chemicals at the following concentrations:

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

<b>Compound</b>	<b>Concentration, mg/l</b>	<b>Compound</b>	<b>Concentration µg/l</b>
NaNO <sub>3</sub>	25.5	H <sub>3</sub> BO <sub>3</sub>	185.52
MgCl <sub>2</sub> .6H <sub>2</sub> O	12.164	MnCl <sub>2</sub> .4H <sub>2</sub> O	415.38
CaCl <sub>2</sub> .2H <sub>2</sub> O	4.41	ZnCl <sub>2</sub>	3.27
MgSO <sub>4</sub> .7H <sub>2</sub> O	14.7	CoCl <sub>2</sub> .6H <sub>2</sub> O	1.428
K <sub>2</sub> HPO <sub>4</sub>	1.044	CuCl <sub>2</sub> .2H <sub>2</sub> O	0.012
NaHCO <sub>3</sub>	*see above	Na <sub>2</sub> MoO <sub>4</sub> .2H <sub>2</sub> O	7.26
		FeCl <sub>3</sub> .6H <sub>2</sub> O	159.76
		Na <sub>2</sub> EDTA.2H <sub>2</sub> O	300.0

*S. capricornutum* cultures were maintained by the testing laboratory at a temperature of 24±2 °C under constant illumination of 4300 10% lux provided by cool-white fluorescent bulbs. The source of the culture was the Department of Botany, University of Texas (initial strain #1648). The algal inoculum used to start the test originated from 5-day old stock cultures in log phase growth.

To start the test, the different WAF and control solutions were placed in their respective number of replicate flasks and inoculated with *S. capricornutum* to an initial cell density of  $1.0 \times 10^3$  cells/ml. The flasks were sealed and randomly positioned on a shaker table. All flasks were incubated for four days under constant illumination (4200 to 4400 Lux), continuous shaking at 125 rpm, and at a mean temperature of 23.6 °C.

Cell density was determined for each replicate flask at 24, 48, 72, and 96 hours. This was done by taking 3.5 ml of solution from each flasks and measuring fluorescence on a Turner Filter fluorometer and converting fluorescence readings to algal cell densities using a regression formula developed through cell counts. In order to maintain no headspace in the test flasks, the 3.5 ml aliquot of solution removed for analyses was replaced with 3.5 ml of spare WAF. The spare WAF samples were stored in closed vessels, and a new vessel was opened for each replacement interval. The replacement of 3.5 ml of WAF diluted the cell densities by approximately 2.5% on Days 2-4. The overall affect of this was considered not significant.

The pH of the control and test solutions were taken at 0 and 96 hours. The pH of the solutions ranged from 7.5 to 7.6 at the beginning of the test and 7.7 to 9.9 at the end of the test.

The EL<sub>50</sub> values were determined on the percent inhibition relative to the control values for average specific growth rate and area under the growth curve. The specific growth rates for each treatment were determined by calculating the slope of the regression line of ln(cell density) versus time. The area under the growth curves was calculated in accordance with the equations in OECD Guideline No. 201.

**Reliability** : (1) valid without restriction

(49)

## 4. Ecotoxicity

Id Kerosene  
Date 9/21/2010

### 4.4 CHRONIC AQUATIC INVERTEBRATE TOXICITY

**Category Chemical :** Kerosene/Jet Fuel category

**Test Substance :** Kerosene (petroleum), hydrodesulfurized

**Test Substance Purity/Composition and Other Test Substance Comments :** The test substance is a complex combination of hydrocarbons obtained from a petroleum stock by treating with hydrogen to convert organic sulfur to hydrogen sulfide which is then removed. It consists of hydrocarbons having carbon numbers predominantly in the range of C9 through C16 and boiling in the range of approximately 150 to 290°C.

**Category Chemical Result Type :** Measured

**Test Substance Result Type:** Measured

#### Method

**Year Study Performed :** 2010

**Method/Guideline Followed:** OECD Guideline 211

**Deviations from Method/Guideline :** Due to the complex mixture of low solubility hydrocarbons in the test substance, the exposure solutions were prepared as individual water accommodated fractions (WAF) in accordance with OECD recommendations.

**Species:** Daphnia magna Straus

**GLP:** Yes

**Analytical Monitoring :** Yes

**Test Type:** Static renewal

**Test Vessel:** Glass screw-top bottles having a capacity of 130 mL without headspace

**Water Media Type:** Reconstituted hard water

**Test Concentrations:** WAFs prepared at the following loading rates: 0 (control), 0.08, 0.19, 0.48, 1.2, and 3.0 mg/L

**Nominal and Measured Concentrations:** Nominal: 0 (control), 0.08, 0.19, 0.48, 1.2, and 3.0 mg/L  
Mean Measured: 0 (<PQL), 0.0163, 0.0392, 0.0916, 0.228, 0.537 mg/L as total dissolved hydrocarbons  
PQL = Practical Quantification Limit = 0.016 mg/L

# 4. Ecotoxicity

**Id** Kerosene  
**Date** 9/21/2010

**Total Exposure Period:** 21 days

<b>Vehicle Used:</b>	None	
<b>Vehicle Name:</b>		
<b>Vehicle Amount and Units:</b>		
<b>Alkalinity:</b>	109 mg/L as CaCO <sub>3</sub>	
<b>Dissolved Oxygen:</b>	5.61 – 9.97	
<b>pH Value:</b>	Value or Lower Range : 8.01 Upper Range : 8.81	
<b>Test Temperature and Units:</b>	Value or Lower Range : Upper Range :	19.8 21.8
<b>Photo (Light/Dark):</b>	16 H light / 8 H dark	
<b>Salinity:</b>	N/A	
<b>TOC:</b>		
<b>Water Hardness:</b>	Value or Lower Range: Upper Range:	160 192



## 4. Ecotoxicity

Id Kerosene  
Date 9/21/2010

Preparation of the exposure solutions followed the WAF procedure. Individual treatments were prepared by adding the test substance to dilution water in glass aspirator bottles and stirring on magnetic stir plates with a vortex of <10% of the static liquid depth for approximately 24 hours. Stirring was stopped after 24 hours and the aqueous portions (water accommodated fractions) of each treatment solution were allowed to settle for approximately 1 hour. Thereafter the WAF solutions were removed and used in testing. The loading rates were 0 (control), 0.08, 0.19, 0.48, 1.2, and 3.0 mg/L. Water quality measurements (pH, dissolved oxygen, hardness, and temperature) were taken from fresh WAF samples and composite samples of old WAF solutions during the study.

**Method/Guideline**  
**Test Conditions**  
**Remarks:**

Ten replicates were established for each test substance treatment and the control using one daphnid per replicate. Replicate chambers were completely filled, with no headspace. Test solutions were renewed every 48 hours with WAFs which had been prepared the previous day. The daphnids were observed daily for immobilization, reproduction, and abnormal behavior/appearance and the numbers of young were counted on transfer days and noted on the days in between transfer days. At the end of the test, the total number of living offspring produced per live parent animal was determined. Adult organisms were measured (body length excluding the anal spine) at termination in order to determine if growth effects occurred.

Daphnids were fed during renewals by adding between 0.325 mL and 0.447 mL of a  $1.3 \times 10^8$  cells/mL suspension of *Pseudokirchneriella subcapitata* to provide approximately  $3.25 \times 10^5$  -  $4.47 \times 10^5$  cells/mL. Test organisms were also fed during renewals with between 0.163 mL and 0.244 mL of a YTC daphnid feed mixture. Feed levels were increased throughout the duration of the study to compensate for daphnid growth and the presence of neonates. The algae and YTC daphnid feed mixture were supplied by Aquatic Biosystems, Inc., Fort Collins, Colorado.

Fresh and old WAF solutions were analyzed for dissolved hydrocarbons by automated static headspace gas chromatography with flame ionization detector (HS GC-FID). Instrumentation was a Pekin-Elmer HS 40 Headspace Sampler connected to a Perkin Elmer AutoSystem XL Gas Chromatograph and FID.

Limit Test: N/A

**Test Results**

**NOEC/LOEC/NOELR/LOELR**

	Exposure Duration:	Exposure Units:	Value Description:	Value or Lower Range:	Upper Range:	Units:	Basis for Concentration:
NOEC:	21	Days	=	0.092		mg/L	Measured
LOEC:	21	Days	=	0.23		mg/L	Measured
NOEL R:	21	Days	=	0.48		mg/L	Nominal loading rate
LOEL R:	21	Days	=	1.2		mg/L	Nominal loading rate

**LC/EC/IC/EL/LL Mean Value**

## 4. Ecotoxicity

Id Kerosene

Date 9/21/2010

Exposure Duration:	Exposure Units:	Type	%:	Value Description:	Mean Value or Lower Mean Value:	Upper Mean Value:	Units :	Basis for Effect:	Basis for Concentration:
21	days	LL	20	=	0.53		mg/L	Reproduction	WAF loading rate
21	days	LL	50	=	0.89		mg/L	Reproduction	WAF loading rate
21	days	LL	20	=	0.41		mg/L	Adult immobility	WAF loading rate
21	days	LL	50	=	0.81		mg/L	Adult immobility	WAF loading rate
21	days	LC	20	=	0.10		mg/L	Reproduction	Measured
21	Days	LC	50	=	0.16		mg/L	Reproduction	Measured
21	days	LC	20	=	0.080		mg/L	Adult immobility	Measured
21	days	LC	50	=	0.15		mg/L	Adult immobility	Measured

### Results Remarks:

The control daphnids released their first brood between days 8 and 12. The coefficient of variation for control fecundity was 9.9%. The test was considered acceptable based on exceeding the minimum control survival (90%) and the reproductive output exceeded the minimum of 60 young per adult. The analytical measurements of the dissolved hydrocarbons in the fresh and old WAF solutions indicated that the exposure concentrations were adequately maintained over the renewal period. Mean concentrations in the old solutions were between 78% and 100% of the initial concentrations.

Prior to the definitive life cycle test, a WAF equilibration and stability evaluation was done. The equilibration/stability trial established that a mixing duration of 24 hours provided sufficient dissolution of the kerosene hydrocarbons into the WAF. Additionally, this trial provided evidence that the hydrocarbons in the WAFs would be adequately stable during the 48-hour intervals between WAF renewals.

### Reliability/Data Quality

Reliability: 1

Reliability Remarks: Reliable without restrictions

Key Study Sponsor Indicator: Key

### Reference

Reference: ExxonMobil Biomedical Sciences, Inc. 2010. *Daphnia magna*, Reproduction Test on Kerosene. Study No. 0950146. ExxonMobil Biomedical Sciences, Inc., Annandale, New Jersey.

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

### 5.1.1 ACUTE ORAL TOXICITY

**Type** : LD<sub>50</sub>  
**Value** : > 5000 mg/kg bw  
**Species** : Rat  
**Strain** : Sprague-Dawley  
**Sex** : Male/female  
**Number of animals** : 5  
**Vehicle** : Undiluted  
**Doses** : Single dose of 5 g/kg bwt  
**Year** : 1985  
**GLP** : Yes  
**Test substance** : Straight run kerosene, sample API 83-09 (See section 1.1.1.)

**Method** : The test material was administered by oral gavage as a single dose of 5 g/kg to five male and five female Sprague-Dawley rats. Food was withheld overnight before administration of test material whilst water was available ad libitum. The animals were observed for clinical signs and mortality at hourly intervals for the first six hours after dosing and twice daily thereafter for 14 days. Body weights were recorded before fasting, just prior to administration of test material and 7 and 14 days after test material administration. At the end of the study, all animals were killed and underwent a gross necropsy. Any abnormalities were recorded.

**Result** : Animals gained weight following administration of the test material. Clinical signs observed included: hypoactivity, ataxia, prostration, soft stool, lacrimation, yellow-stained abdomen and/or urogenital region and hair loss on abdomen and/or urogenital region. There were no mortalities during the study. There were no gross abnormalities at necropsy.

**Reliability** : (1) valid without restriction (16)

**Type** : LD<sub>50</sub>

**Result** : Two additional acute oral toxicity studies have been reported for substances in this group. The results were as follows:

<u>Sample</u>	<u>LD<sub>50</sub></u>	<u>Report reference</u>
Hydrodesulfurized kerosene API 81-07	> 5g/kg	API 30-31986
Jet Fuel A	>20 g/kg	API 27-32815

(8) (10)

### 5.1.2 ACUTE INHALATION TOXICITY

**Type** : LC<sub>50</sub>  
**Value** : > 5 mg/l  
**Species** : Rat  
**Strain** : Sprague-Dawley  
**Sex** : male/female  
**Number of animals** : 5  
**Vehicle** : Air

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

- Doses** : 5.28 ±0.42 mg/l  
**Exposure time** : 4 hour(s)  
**Year** : 1987  
**GLP** : Yes  
**Test substance** : Straight run kerosene, sample API 83-09 (See section 1.1.1.)
- Method** : Five male and five female Sprague-Dawley rats were exposed to approximately 5 mg/l of test material as a single four-hour whole body inhalation exposure. After the exposure the rats were kept for a 14 day observation period. Surviving animals at 14 days were sacrificed and subjected to a gross post-mortem examination. Records were made of any observed gross abnormalities. The lungs of all animals were preserved in formalin, sectioned and stained and then subjected to microscopic examination.  
Since no animals died following the exposure, no further exposures were conducted.
- Result** : All except one animal had normal growth rates throughout the study. The one exception on day 8 had a body weight less than its starting body weight but by the end of the study normal growth had resumed. Decreased activity was exhibited by all animals during the exposure. Otherwise there were no treatment-related clinical signs of toxicity.  
No macroscopic lesions were observed in any animal at post-mortem and no microscopic changes were observed in any lung section examined.
- Test condition** : Exposure of the animals was conducted in a 250 liter stainless steel and glass chamber.  
The test material was introduced at a rate of 0.34 to 0.51 ml/minute into the top of a vertical counter current column that was packed with steel mesh and heated to approximately 50 °C. Pre-warmed nitrogen gas was introduced at the bottom of the column at a rate of 7 liters/minute. The nitrogen and test material vapors were then mixed with air and introduced into the exposure chamber. The test atmosphere was analyzed by IR every 15 minutes throughout the exposure.  
During each hour of the exposure, the test atmosphere was analyzed gravimetrically and visually (by flashlight) for aerosols of the test material. Test material consumption was determined by weighing the test material container before and after exposure. The quantity consumed was divided by the total airflow and this yielded the nominal exposure concentration.  
The mean exposure concentrations were:
- |                          |                 |
|--------------------------|-----------------|
| Nominal concentration    | 5.74 mg/l       |
| Analytical concentration | 5.28 ±0.42 mg/l |
- Gravimetric samples, collected on membrane filters, and aerosol checks with a flashlight showed some aerosol in the chamber. The nominal to analytical ratio and the gravimetric results both suggest the level of aerosol compared to level of vapor was insignificant in the exposure.
- Reliability** : (1) valid without restriction

(21)

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

**Type** : LC<sub>50</sub>  
**Exposure time** :  
**Result** : Additional acute inhalation toxicity studies have also been reported for substances in this group.

<u>Sample</u>	<u>LC<sub>50</sub></u>	<u>Report reference</u>
(Species) Hydrodesulfurized kerosene API 81-07 (Rat, 4-hour)	> 5.2 mg/l	API 30-32855
Deodorized kerosene (Rat, 8-hour)	No mortalities when exposed to saturated vapor	Carpenter et al
(Cat, 6-hour)	>6.4 mg/l	Carpenter et al (12) (34) (43)

### 5.1.3 ACUTE DERMAL TOXICITY

**Type** : LD<sub>50</sub>  
**Value** : > 2000 mg/kg bw  
**Species** : Rabbit  
**Strain** : New Zealand white  
**Sex** : Male/female  
**Number of animals** : 2  
**Vehicle** : Undiluted  
**Doses** : 2 g/kg  
**Year** : 1985  
**GLP** : Yes  
**Test substance** : Straight run kerosene, sample API 83-09 (See section 1.1.1.)

**Method** : Undiluted test material was applied to the shorn dorsal skin of each of two male and two female rabbits. One rabbit of each sex had abraded skin, the other had intact skin. The area of application was wrapped with gauze and overwrapped with an occlusive covering. 24 hours later, the covering was removed and the skin was wiped with wet disposable towels to remove any residual test material.

The rabbits were observed for clinical signs and mortality for the first six hours of dosing, then daily for dermal irritation and twice daily for clinical signs of toxicity and mortality for 14 days. Body weights were recorded just prior to and again 7 days and 14 days after administration of test material. At study termination all animals were killed and subjected to a gross necropsy examination when any abnormalities were recorded.

**Result** : Clinical signs observed during the study included hypoactivity and diarrhea. Dermal irritation ranged from  
- slight to severe for erythema and edema  
- slight to marked for atonia, desquamation and fissuring  
- slight to moderate for coriaceousness.  
Other dermal irritation observed included subcutaneous hemorrhage, blanching and scab formation.  
No animals died during the study.  
One of the rabbits (male abraded skin) weighed slightly less at the end of

## 5. Toxicity

Id Kerosene

Date 9/21/2010

- the study than at the beginning. All other rabbits had gained a small amount of body weight by the end of the study.
- Reliability** : (1) valid without restriction (16)
- Type** : LD<sub>50</sub>
- Result** : Two additional acute dermal toxicity studies have been reported for substances in this group. The results were as follows:
- | <u>Sample</u>                        | <u>LD<sub>50</sub></u> | <u>Report reference</u> |
|--------------------------------------|------------------------|-------------------------|
| Hydrodesulfurized kerosene API 81-07 | > 2g/kg                | API 30-31986            |
| Jet Fuel A                           | >4 g/kg                | API 27-32815            |
- (8) (10)

### 5.2.1 SKIN IRRITATION

- Species** : Rabbit
- Concentration** : Undiluted
- Exposure** : Occlusive
- Exposure time** : 24 hour(s)
- Number of animals** : 6
- Vehicle** : None
- PDII** : 5.5
- Year** : 1985
- GLP** : Yes
- Test substance** : Straight run kerosene, sample API 83-09 (See section 1.1.1.)

- Method** : Six rabbits were used in this study. Approximately 24 hours before the study the hair was clipped from the back and flanks of each animal. Just prior to the test material administration, abraded areas were prepared; these were not sufficiently deep to cause bleeding. 0.5 ml of test material was applied to two areas of skin on each animal (abraded and intact skin). The treated areas were each covered with a gauze which was secured with paper tape and then covered with an occlusive dressing. After 24 hours exposure, the patches were removed and residual test material was removed from the skin by gentle wiping with a wet paper towel. The degree of erythema and edema was recorded using the Draize scale. A second assessment of skin reaction was made at 72 hours. Because irritation was still present at 72 hours, further evaluations of skin irritation were made at 96 hours, 7 days and 14 days post exposure. Body weights were recorded just prior to the application of the test material and again at weekly intervals throughout the study.

- Result** : The results are given in the following table.

<b>Observation time</b>	<b>Erythema</b>		<b>Edema</b>		<b>Total score</b>
	<b>Intact</b>	<b>Abraded</b>	<b>Intact</b>	<b>Abraded</b>	
24 hrs	2.5	2.8	2.8	3.2	5.7
72 hrs	2.8	3.0	2.3	2.2	5.2
96 hrs	2.8	3.0	2.7	2.5	5.5
7 days	2.3	2.3	2.0	2.2	4.4
14 days	0.3	0.7	0.0	0.0	0.5

Primary dermal irritation Index= 5.5

The primary dermal irritation index is the sum of the irritation scores for 24

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

and 72 hours divided by 2 and rounded to the nearest tenth.  
The test material produced moderate to severe levels of irritation.  
Blanching was seen within the test sites of two animals at 24 and 72 hours and of three animals at 96 hours. Subcutaneous hemorrhages were also observed in two animals at 96 hours.

There were no signs of systemic toxicity.  
**Reliability** : (1) valid without restriction (16)

**Species** : Rabbit

**Result** : A further six skin irritation studies have been reported for this group of substances. The results are:

<b>Material</b>	<b>Skin irritation</b>	<b>Reference</b>
Hydrodesulphurized kerosene API 81-09	Mild-moderate (24 hr)	API 30-31986
Odorless kerosene	Mild (4 hour)	Shell SBER 91-001
Kerosene SG	Mild ( 4 hour)	Shell SBER 91-003
Hydrocracked kerosene	Mild (4 hour)	Shell SBER 91-004
Jet A-1	Mild (4 hour)	Shell 91-005
Jet fuel A	Moderate-severe (24 hr)	API 27-32815 (8) (10) (65) (66) (67) (68)

### 5.2.2 EYE IRRITATION

**Species** : Rabbit  
**Concentration** : Undiluted  
**Dose** : 0.1 ml  
**Comment** : One group of animals, eyes rinsed after 30 seconds  
**Vehicle** : None  
**Year** : 1985  
**GLP** : Yes  
**Test substance** : Straight run kerosene, sample API 83-09 (See section 1.1.1.)

**Method** : 0.1 ml undiluted test material was dripped onto the corneal surface of one eye of each of nine rabbits. The upper and lower eyelids were held together for one second to prevent loss of test material. 20-30 seconds after application of test material the eyes of three rabbits were flushed for 1 minute with lukewarm water. The other six rabbits did not receive any further treatment.

The eyes of each rabbit were examined 1, 24, 48 and 72 hours and seven days after application of test material. Sodium fluorescein was as an aid to reveal possible corneal injury. Grading and scoring of ocular lesions was performed according to the Draize scale. Body weights were recorded just before treatment and again at the end of the study.

**Result** : A pain response was elicited in one animal following instillation of test material. No corneal or iridial irritation was seen during the study. All irritation had cleared by the 24 hour observation time.  
No systemic toxicity was seen during the study and body weights were unaffected by treatment. The primary irritation score recorded at the 1 hour observation period was:

## 5. Toxicity

Id Kerosene

Date 9/21/2010

<u>Unwashed eyes</u>	<u>Washed eyes</u>
----------------------	--------------------

6 rabbits	3 rabbits
-----------	-----------

1 hour 0.7	2.0
------------	-----

The scores for all other observation times were zero.

The primary irritation score is the total eye irritation score for all animals, divided by the number of animals in each group.

**Reliability** : (1) valid without restriction (16)

**Species** : Rabbit

**Result** : Two other eye irritation studies have been reported.

For these studies on hydrodesulphurized kerosene (API Report No. 30-31986) and jet fuel (API report 27-32815), the method of scoring ocular lesions and calculating primary eye irritation index was exactly comparable to the study described above on straight run kerosene. The results for these two materials are as follows:

Test material	Primary eye irritation index	
	Unwashed eyes	Washed eyes

Hydrodesulphurized kerosene

1 hour	3.0	2.7
--------	-----	-----

24 hours	0.3	0.7
----------	-----	-----

48 hours	0.3	0
----------	-----	---

72 hours	0	0
----------	---	---

7 days	0	0
--------	---	---

Jet fuel

24 hour	3.33	1.33
---------	------	------

48 hour	1.0	1.33
---------	-----	------

72 hour	1.0	1.33
---------	-----	------

7 day	0	0
-------	---	---

(8) (10) (42)

### 5.3 SENSITIZATION

**Type** : Buehler Test  
**Species** : Guinea pig  
**Concentration** : 1<sup>st</sup>: Induction 75 % occlusive epicutaneous  
2<sup>nd</sup>: Challenge 10 % occlusive epicutaneous  
**Number of animals** : 10  
**Vehicle** : Paraffin oil  
**Result** : Not sensitizing  
**Year** : 1985  
**GLP** : Yes  
**Test substance** : Straight run kerosene, sample API 83-09 (See section 1.1.1.)

**Method** : 0.4 ml undiluted test material was applied under an occlusive dressing to the shaved skin of 10 animals. Six hours after application, the dressing was removed and the skin wiped to remove residues of test material. The animals received one application each week for 3 weeks. The same skin site was used for the first two applications but due to the severe irritation that had occurred in the test and positive control groups a different site was used for the third sensitizing application. 2 weeks following the third application a challenge dose (0.4 ml of a 1% solution in paraffin oil) was applied in the same manner as the sensitizing



## 5. Toxicity

Id Kerosene  
Date 9/21/2010

doses. A previously untreated site was used for the challenge application. The application sites for sensitizing and challenge doses were scored for erythema and edema 24 and 48 hours after patch removal. To assist in the scoring of the response to the final challenge dose the test site was depilated 3 hours prior to examination by using a commercially available depilatory cream.

Positive control (2,4-dinitrochlorobenzene), vehicle control and naive control groups were included in this study.

Concentrations of positive control were as follows:

Sensitizing doses: 0.4 ml of 0.3% w/v in 80% aqueous ethanol

Challenge dose: 0.4 ml of 0.1% w/v suspension in acetone

**Result** : The skin reactions after challenge applications were as follows:

test group No dermal irritation in any animal

Naive control group Very slight erythema in two animals, no reaction in the other eight animals

Vehicle control group A very slight erythema in one animal, no reaction in the other nine animals

Positive control group Very slight to severe irritation in all 20 animals. The reaction in 19 animals exceeded the highest reaction observed in the naive positive control animals

Naive positive control group 5 of 20 animals exhibited very slight erythema, the other 15 animals had no skin reaction.

**Reliability** : (1) valid without restriction

(16)

**Type** : Buehler Test

**Result** : Jet Fuel A was not a skin sensitizer in a Beuhler test (Report API 27-32815).

Hydrodesulfurized kerosene was not a skin sensitizer in a Beuhler test (Report API 31-31413).

(8) (13)

### 5.4 REPEATED DOSE TOXICITY

**Type** :  
**Species** : Rabbit  
**Sex** : Male/female  
**Strain** : New Zealand white  
**Route of admin.** : Dermal  
**Exposure period** : 28 days  
**Frequency of treatm.** : Three times weekly  
**Doses** : 200, 1000 and 2000mg/kg/day  
**Control group** : Yes, concurrent no treatment  
**Method** :  
**Year** : 1985  
**GLP** : No  
**Test substance** : Straight run kerosene, sample API 83-09 (See section 1.1.1.)

**Method** : Undiluted test material was applied to the shorn dorsal skin of each of five male and five female rabbits at doses of 200, 1000 and 2000 mg/kg/day,

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

three times weekly until 12 doses had been applied. Five rabbits of each sex served as sham treated controls. Dosing was carried out on alternate days.

Each treated site was covered with a gauze pad and an occlusive dressing. The occluded dressing was then removed after six hours and any residual test material was removed from the skin with a clean, dry, absorbent gauze pad.

Each animal was observed twice daily for clinical signs of toxicity and pharmacological effects. Body weights were recorded prior to dosing and then weekly throughout the study. The test site was examined daily and dermal reactions were graded using the Draize scale and recorded. At termination, blood samples were collected for the following clinical chemical and hematological determinations.

<b>Hematology</b>	<b>Clinical chemistry</b>
Erythrocyte count	Glucose
Total leukocyte count	Blood urea nitrogen
Differential leukocyte count	Total protein
Hemoglobin	SGOT
Hematocrit	SGPT

All animals were sacrificed and necropsied whether they had died or had survived throughout the study. Organs from animals found dead were not weighed but for animals surviving to the end of the study, the following organs were weighed and organ/body weight ratios were determined. Heart, liver, spleen, kidneys, adrenals, thyroid, pituitary, testes, ovaries and brain.

The following tissues were collected and preserved and were prepared for subsequent histological examination.

Heart, Sacculus rotundus, Urinary bladder, lungs, colon, adipose tissue, bronchi, thymus, mammary gland, trachea, spleen, brain (cerebellum, cerebrum, pons), thyroid, liver, parathyroids, pancreas, pituitary, cervical lymph nodes, kidneys, spinal cord (two sections), salivary gland, adrenals, skeletal muscle, tongue, vagina, sciatic nerve, esophagus, seminal vesicles, testes/ovaries, skin (treated and untreated), stomach, bone, duodenum, epididymis, bone marrow (smear), jejunum, ileum, prostate/uterus, eyes, mesenteric lymph nodes and any gross lesions.

A two-tailed student's t-test was used to determine the significance of any differences between treated and control groups for body weights, clinical pathology and absolute and relative organ weights.

**Result** : One control male was found in a moribund state and was sacrificed on day 21 of the study. One female control was found dead on day 11 of the study.  
One 1000 mg/kg/day male was found dead on day 15.  
A male and a female in the highest dose group were found dead on days 10 and 24 respectively and the authors considered these to be treatment-related.  
Clinical signs observed in the study that were considered to be treatment-related included: thinness, nasal discharge, lethargy, soiled anal area, anal discharge, wheezing.  
There were group mean body weight losses in the mid and high dose groups and a smaller increase than controls in the low dose group as follows:

## 5. Toxicity

Id Kerosene

Date 9/21/2010

Group	Weight gain (kg)	
	Males	Females
Control	0.3	0.3
200 mg/kg/day	0.2	0.2
1000 mg/kg/day	0	0
2000 mg/kg/day	-0.3*	-0.4*

P < 0.05

The authors judged that the weight losses in the mid dose group were not a direct effect of the test material. They noted that weight losses occurred in the first week of dosing and that after this there were increases in weight. The authors comment that such effects generally occur as a result of the stress of dosing and the dermal irritation that occurred.

The skin irritation grades showed that irritation was dose related and was greatest in the highest dose group. The mean irritation score, calculated as the mean sum of all the irritation scores (erythema and edema), for each group was as follows:

Group/sex	MIS	Classification
Control Male	0	Non-irritant
Control Female	0	Non-irritant
200 mg/kg/day Male	1.3	Slight irritant
200 mg/kg/day Female	1.7	Slight irritant
1000 mg/kg/day Male	3.7	Moderate irritant
1000 mg/kg/day Female	3.5	Moderate irritant
2000 mg/kg/day Male	4.1	Moderate irritant
2000 mg/kg/day Female	3.6	Moderate irritant

Other dermal findings included cracked, flaky and/or leathery skin, crusts and/or hair loss. These findings only occurred in the treated groups and appeared with greater frequency as the dose level increased.

There were no hematological findings in the female groups.

In males, reductions in RBC, hemoglobin and hematocrit were as shown. The actual mean values are shown for the controls and the % reductions are shown for the treated groups. Those values indicated \* are significantly different (P<0.05).

	Control	200 mg/kg	1000 mg/kg	2000 mg/kg
RBC	7.1	-13%*	-20%*	-11%
Hemoglobin	14.9	-6%*	-15%*	-15%*
Hematocrit	45.2	-7.7%*	-21%*	-16%*

No treatment-related changes were seen in the clinical chemistry data with the following exceptions

Total protein reduced by 8% in 200 mg/kg males; SGPT and ALP reduced by 37% and 46% respectively in the 2000 mg/kg females. The authors comment that these values were well within the normal range of historical controls and are not judged to be test material related.

Organ weights and organ/body weight ratios differed from controls as shown in the following table. Values are shown as % differences, either greater (+%) or less (-%) than corresponding controls. Note that all values have been rounded to nearest whole number and only those values significantly different from controls are shown. No differences were observed for any other organ weight.

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

Organ	Dose mg/kg	Absolute organ wt		Relative organ wt	
		M	F	M	F
Heart	200	+17%		+18%	
	1000	+22%		+32%	+35%
	2000				+35%
Liver	200				
	1000				
	2000	-24%			
Spleen	200		+128%		+125%
	1000	+83%	+91%	+75%	+125%
	2000			+50%	
R Kidney	200				
	1000				+20%
	2000			+13%	+28%
L Kidney	200				
	1000				+15%
	2000			+60%	+19%
R Adrenal	200		+55%		+50%
	1000				
	2000		+67%		+100%
L Adrenal	200				+75%
	1000				
	2000		+54%		+100%
Pituitary	200				
	1000				
	2000		+21%		+100%
Thyroid	200				
	1000				
	2000				
Brain	200				
	1000				
	2000			+21%	+21%
R Testis	200				
	1000				
	2000				
L Testis	200				
	1000				
	2000				
R Ovary	200				
	1000				
	2000				
L Ovary	200				
	1000				
	2000				

The authors concluded that the increases in relative heart weights for the mid- and high- dose males and females were treatment-related. Other heart weight changes were within the normal range for control values for the laboratory. Increased absolute and relative spleen weights for males were considered incidental since they fell within the normal range for the laboratory. For the females however, the differences were considered to be treatment-related. In both males and females, differences in absolute and relative adrenal weights were considered to be stress-related and therefore, indirectly related to treatment.

Gross necropsy findings were confined largely to the skin. Enlarged

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

spleens in the female groups were also noted.

### Microscopic pathology

Slight to moderate proliferative and slight to moderately severe inflammatory changes were present in the treated skin of all male and female animals in the high dose group. These changes were accompanied by an increase in granulopoiesis of the bone marrow in 5/6 males and 3/4 females.

4/6 high dose group males also had multifocal or diffuse tubular hypoplasia of a few of the seminiferous tubules of both testes. The degree of spermatogenesis was similar to controls in one animal, was absent in two animals and was slightly reduced in three animals. These testicular changes were considered by the authors to be secondary to the skin and/or weight changes.

All other lesions observed were considered to be incidental and unrelated to treatment.

**Reliability** : (1) valid without restriction  
This study was audited by a quality assurance unit and was found to be satisfactorily conducted and reported.

(15)

**Type** : Sub-chronic  
**Species** : Rat  
**Sex** : Male/female  
**Strain** : Sprague-Dawley  
**Route of admin.** : Dermal  
**Exposure period** : Six hours each day  
**Frequency of treatm.** : Daily, five days per week for 13 weeks  
**Doses** : 165, 330 & 495 mg/kg/day  
**Control group** : Yes, concurrent vehicle  
**Year** : 1997  
**GLP** : Yes  
**Test substance** : Hydrodesulfurized kerosene

**Method** : Groups of 12 male and 12 female, individually housed, Sprague-Dawley rats aged 7-9 weeks were used. The males weighed 198-328 g and the females weighed 156-249 g at the initiation of the study. Test material was applied at concentrations of 20, 40 or 60% (v/v) at a rate of 1 ml/kg/day to the shorn intrascapular region of the rats. This was equivalent to doses of test material of 165, 330 or 495 mg/kg/day. Dosing was continued daily for five consecutive days each week, five days a week for 13 weeks. In addition a group of 12 male and 12 female rats of similar age were administered mineral oil at a dose rate of 1 ml/kg/day; these animals served as vehicle controls. An additional 12 rats/sex/group in the vehicle controls and high dose group were maintained for a 4-week recovery period following dosing for 13 weeks. All animals were fitted with collars to prevent ingestion and these were removed six hours after dosing and any residual test or control material was wiped from the skin. Animals were observed for clinical signs prior to dosing and 1, 6 and 24 hours after the first dose. Subsequently, observations were made prior to each dose being applied. Prior to the administration of each dose, the treated skin site was evaluated for dermal irritation using the Draize scoring method. Body weights were recorded prior to the first dose and weekly thereafter. An ophthalmic examination was conducted on each rat prior to application of the first dose and again prior to sacrifice at the end of the study. During the week prior to the first dose, each rat was subjected to a

functional observation battery (FOB). The FOB was conducted again 1, 6 and 24 hours after the first dose and at 7 and 14 days. During the study, the FOB, motor activity and startle response testing was conducted on all rats at weeks 4, 8 and 12.

[The details of the FOB, the test for startle response test and the test for motor activity are given in detail in the laboratory report but are not included here].

At the 14 week necropsy, blood samples were collected from 12 animals/sex/group and at the week 18 necropsy from the recovery rats (vehicle and high dose groups).

The following hematological and clinical chemical parameters were measured.

### **Hematology**

Erythrocyte count

Hemoglobin

Hematocrit

Mean corpuscular volume

Mean corpuscular hemoglobin

Mean corpuscular hemoglobin concentration

Platelet count

Reticulocyte count

Total leukocyte count

Differential leukocyte count

Morphological examination of erythrocytes and platelets

Coagulation determinations (prothrombin time & activated partial thromboplastin time) were also carried out on six animals from each group at week 14 and from the recovery groups at the week 18 necropsy.

### **Clinical chemistry**

Blood urea nitrogen

Creatinine

Serum aspartate aminotransferase

Serum alanine aminotransferase

Alkaline phosphatase

Lactate dehydrogenase

Sorbitol dehydrogenase

Gamma glutamyl transferase

Creatinine kinase

Serum glucose

Total, direct and indirect bilirubin

Total protein

Albumin

Calcium

Phosphorus

Sodium

Potassium

Chloride

A complete necropsy was performed on six rats/sex/group following 13 weeks dosing, and on 6 rats/sex/group of the recovery animals (high dose and controls) at week 18. A limited necropsy was performed on the remaining six animals and their organs were not weighed (see below).

Each full necropsy included an examination of the external surface of the body, all orifices, cranial, thoracic, abdominal and pelvic cavities and their contents. Gross observations were recorded and the following organs were weighed:

## 5. Toxicity

Id Kerosene

Date 9/21/2010

Adrenal glands, brain, epididymides, heart, kidneys, liver, ovaries, prostate, spleen, testes, thymus and uterus.

The following tissues were collected, processed and then examined microscopically.

Adrenal glands	Nose (nasal cavity & turbinates)
Animal identification	Ovaries
Bone marrow (from sternum)	Oviducts
Brain	Pancreas
Epididymides	Parathyroid glands
Esophagus	Pituitary gland
Exorbital lacrimal glands	Prostate
Eyes with optic nerve	Salivary glands
Femur (incl. articular surface)	Seminal vesicles
Gross lesions	Skin (application site)
Harderian gland	Skin (inguinal)
Heart and aorta	Spinal cord (3 levels)
Intestine (3 levels)	Spleen
Kidneys	Stomach
Larynx and pharynx	Testes
Liver	Thymus
Lungs with mainstream bronchi	Thyroid gland
Lymph nodes (mandibular/mesenteric)	Urinary bladder
Mammary glands with adjacent skin	Uterus
Muscle (thigh)	Vagina
Nerve (sciatic)	

The remaining six rats of each group were anesthetized with an intraperitoneal injection of Pentothal ® and transcardially perfused in-situ using 10% neutral-buffered formalin and given a limited necropsy. For these rats, no organs were weighed and the following tissues were collected:

Head/skull	Sural nerve
Brain	Tibial nerve
Spinal cord	Gross lesions
Sciatic nerve	

The following tissues were examined microscopically in these animals:  
Brain (forebrain, cerebrum, midbrain, cerebellum, pons and medulla obligata)

Gasserian ganglia  
Dorsal root ganglia  
Dorsal and ventral root fibers  
Sural nerve  
Tibial nerve  
Spinal cord (cervical and lumbar areas)  
Sciatic nerve.

### Statistics

Normally-distributed in-life data (parametric) were analyzed for test substance effects by analysis of variance and pairwise comparisons made between groups using Dunnett's test. Nonparametric data (nonhomogenous as determined by Bartlett's) were analyzed using a modified t-test. Statistical significance was reported at the  $P < 0.05$  level. Statistical analyses of neurobehavior data (FOB and motor activity) are described in the results section.

### **Result**

: All animals survived until scheduled termination.  
There were no test substance-related effects on survival, clinical

## 5. Toxicity

Id Kerosene

Date 9/21/2010

observations (apart from skin irritation), neurobehavioral signs or ophthalmological findings. The only clinical observations during the study were related to skin irritation at the application site. There was a generally dose-related increase in the incidence and severity of erythema, edema, epidermal scaling, scab formation, thickening of the skin and ulceration at the treated site. Males seemed to be more sensitive than females. The FOB screen did not demonstrate any substance-related effects. The areas monitored were: behavioral parameters, including autonomic, muscle tone and equilibrium, sensorimotor responses, central nervous system. In addition the test substance had little effect on motor activity or startle response.

Growth rates were unaffected by treatment.

At necropsy no substance-related observations were made for males in any group. In the females there was a suggestion of a possible treatment-related effect which occurred in 7 rats across all groups and consisted of skin crusts or ulceration at the site of application of test material. Hematological and serum clinical parameters were unaffected by treatment.

The only organ weight effects noted were an increase in spleen/body weight and spleen/brain weight ratios in the high dose group females at the 13 week necropsy and an increase in absolute spleen weight in the same dose group females after the 4 weeks recovery period. Since there were no associated microscopic or clinical chemical findings, these differences were not considered to be of biological relevance.

There were no treatment-related microscopic changes in the tissues examined with the exception of the findings in the skin. The skin observations were minimal in nature with a severity score less than 1 on a 1 [low] to 4 [severe] scale. The findings included acanthosis, ulceration, parakeratosis, chronic active inflammation and hyperkeratosis. The males were affected at all doses, however, the effects indicated very little irritation. Recovery group animals revealed complete recovery in the females and minimal hyperkeratosis in the high dose group males. No effects were found in the animals subjected to a detailed neuropathological examination.

### Test substance

: The Hydrodesulfurized kerosene had the following properties.

Boiling point	148.9 °C (300 °F)
Specific gravity	0.825 @ 60 °F
Melting point	Not applicable
% volatile	100
Vapor pressure	0.4 mm Hg @ 68 °F
Evaporation rate (water = 1)	Slower
Vapor density (air = 1)	4.7
Viscosity	1.3 - 2.2 cSt @ 100 °F
% solubility in water	Negligible
Pour point	-34.4 °C (-30 °F)
pH	Not determined
Appearance/odor	Clear liquid with hydrocarbon odor

The vehicle used was Squibb mineral oil.

For dosing, mixtures of hydrodesulfurized kerosene were prepared in the mineral oil at concentrations of 20, 40 and 60% (v/v)

### Reliability

: (1) valid without restriction

(31)



## 5. Toxicity

Id Kerosene

Date 9/21/2010

**Route of admin.** : Dermal

**Result** : Five additional dermal repeat dose studies have been reported for materials in this group. The results are summarized below. It should be noted that the 13 week study was used only as a dose range finding study and did not provide information on systemic toxicity, since it was designed to assess the suitability of the dosing regime with respect to skin irritation.

### Study 1

Test material: Hydrodesulfurized kerosene  
Species: Rabbit  
Duration: 28 days  
Dosing regime: 3x/week  
Doses: 200, 1000 and 2000 mg/kg/day  
Findings: Skin irritation only. Other findings were either incidental or secondary to skin irritation or reduced body weight gain.  
Reference: API 30-32297  
Reliability: 1

### Study 2

Test material: Jet Fuel A  
Species: Rabbit  
Duration: 14 days  
Dosing regime: 5x/week  
Doses: 6.4 g/kg/day  
Findings: Severe skin damage at the treatment areas. Depression and weight loss associated with anorexia. Tissue damage in liver (mottled necrosis and centrilobular degeneration), kidney and bladder (hyperplasia) considered to be secondary to severe skin irritation  
Reference: API 27-32815  
Reliability: 1

### Study 3

Test material: Hydrotreated straight run kerosene  
Species: Mouse  
Duration: 3 weeks  
Dosing regime: 3x/week  
Doses: Dose not specified  
Findings: Degenerative skin changes including necrosis and hyperplasia. The effects were well advanced after one week.  
Reference: Data summarized in CONCAWE 91/51  
Reliability: 4, Reported as a review and without sufficient experimental detail.

### Study 4

Test material: 3 kerosene samples: two hydrotreated, straight run kerosenes and a blend of 70% hydrocracked kerosene and 30% hydrotreated straight run kerosene  
Species: Mouse  
Duration: 1 weeks  
Dosing regime: 3 applications in one week

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

Doses: Dose not specified  
Findings: Minimal macroscopic changes. Earliest changes were inflammation and necrosis of the hair follicles with subsequent degeneration at the skin surface  
Reference: Data summarized in CONCAWE 91/51  
Reliability: 4, Reported as a review and without sufficient experimental detail.

### Study 5

Test material: 2 kerosene samples: a hydrotreated, straight run kerosenes and a blend of 70% hydrocracked kerosene and 30% hydrotreated straight run kerosene  
Species: Mouse  
Duration: 13 weeks  
Dosing regime: Various regimes. Purpose was to determine a dosing regime that would be suitable for a long-term study.  
Doses: 100 or 50 µl/application  
Findings: Samples were highly irritating when 50 µl applied undiluted twice/week. Slightly less irritation occurred when 100 µl applied once/week. A 25% solution in white oil applied at a dose of 50 µl, twice/week was non-irritant. At necropsy no treatment-related effects were observed except those occurring in the skin.  
Reference: Data summarized in CONCAWE 93/55  
Reliability: 4, Reported as a review and without sufficient experimental detail.

(8) (11) (37) (38)

**Type** : Sub-chronic  
**Species** : Rat  
**Sex** : Male/female  
**Strain** : Sprague-Dawley  
**Route of admin.** : Inhalation  
**Exposure period** : Four weeks  
**Frequency of treatm.** : Six hours/day, five days/week for four consecutive weeks  
**Doses** : Nominal: 32 mg/m<sup>3</sup>. Actual: 24 mg/m<sup>3</sup>  
**Control group** : Yes  
**Year** : 1986  
**GLP** : yes  
**Test substance** : API 81-09 (Hydrodesulfurized kerosine)

**Method** : Groups of 20 male and 20 female Sprague Dawley rats (approximately six weeks old) were exposed to a nominal concentration of 25mg/m<sup>3</sup> kerosene by inhalation. Exposures were for approximately six hours each day, five days each week for four consecutive weeks. Control groups of 20 male and 20 females were exposed to filtered air. Animals were observed twice daily for overt signs of toxicity and they underwent a detailed examination once weekly. Body weights were also recorded weekly. At study termination, the animals were killed and blood samples were taken for the following clinical chemical and hematological investigations:

**Hematology**

**Clinical chemistry**

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

Hematocrit	Aspartate aminotransferase
Hemoglobin	Alanine aminotransferase
Erythrocyte count	Alkaline phosphatase
MCH	Glucose
MCV	Urea nitrogen
MCHC	Total protein
Leukocyte count	
Platelet count	
Reticulocyte count	

For all rats, the following organs were weighed and the organ body weight ratios were calculated :

Heart, lung and trachea, liver, kidneys, brain, spleen, adrenals, thyroid/parathyroid, pituitary, testes and ovaries.

The following tissues were removed and preserved:

Adrenals (2)	Aorta
Bone marrow (femur)	Bone marrow smear
Brain (3 levels)	Eye with contiguous Harderian gland
Esophagus	Stomach
Duodenum	Jejunum
Ileum	Cecum
Colon	Rectum
Gonads	Ovary (2)
Kidney (2)	Testis with epididymis (2)
Heart	Liver (3 sections)
Nasal tissues	Lung & trachea (all lobes)
Abdominal lymph nodes	Thoracic lymph nodes
Mammary region lymph nodes	
Pancreas	Pituitary
Sciatic nerve	Prostate & seminal vesicle
Skeletal muscle (thigh)	Skin
Salivary gland (mandibular with submandibular lymph node)	
Spinal cord (cervical, midthoracic & lumbar)	
Spleen	Thymic region
Thyroid/parathyroid complex	
Urinary bladder	Uterus (2 horns & cervix)
Vagina	

The following tissues were examined microscopically in all rats:

Adrenal (2), brain (3 levels: fore, mid & hind), bronchi, esophagus, eye (2), heart, kidney (2), liver, lungs (2), lymph node (mediastinal), ovary (2), pancreas, pituitary, prostate, salivary gland, skin, spleen, stomach, testis (2), thymus, thyroid/parathyroid, trachea, urinary bladder, uterus, all gross lesions.

Statistical analysis

Body weight, hematology, clinical chemistry and organ weight data were analyzed by analysis of variance and Bartlett's test.

Treatment groups were compared to control by sex, using the appropriate t-statistic.

Data containing inequalities or where group variances were heterogeneous were compared using a non parametric approach, by transforming the data into ranks prior to analysis as described by Conover and Iman.

### Result

: There were no treatment-related effects on clinical condition, growth rate organ or organ body weight ratios or on any of the hematological or clinical chemistry determinations.  
Furthermore, there were no treatment-related microscopic changes

## 5. Toxicity

Id Kerosene

Date 9/21/2010

**Test condition** : observed in any of the organs examined.  
 : Atmospheres were generated by atomizing the test material into an atomization chamber. The resulting vapors/aerosols were directed to the chamber inlet where dilution with chamber ventilation air reduced the concentration to the desired level.  
 Nominal concentrations were calculated from test material use rates. Actual concentrations were determined at approximately hourly intervals using a Total Hydrocarbon analyzer.

The nominal and actual concentrations for the study were:

Desired concentration (mg/m <sup>3</sup> )	Exposure concentration (mg/m <sup>3</sup> )			
	Nominal Mean	SD	Actual Mean	SD
25	32	1.74	24	1.61

**Reliability** : (1) valid without restriction

(20)

### 5.5 GENETIC TOXICITY 'IN VITRO'

**Type** : In vitro studies

**Remark** : The In vitro genotoxic potential of kerosenes and jet fuel has been studied in several studies and these are tabulated below. Only one study for each test system is described in detail in the following section on in vitro genotoxicity studies and these are indicated (\*) in the table. For completeness, the results of all studies are briefly summarized as separate robust summaries. At the end of this section the results of all in vitro studies are summarized in tabular form.

Study type	Test material	Reference
<b>Ames assay</b>		
	Deodorized kerosene	API 26-60103
	Straight run kerosene	API 26-60017*
	Jet Fuel A	API 27-30051
<b>Modified Ames assay</b>		
	Straight run kerosene	CONCAWE 91/51*
	Hydrotreated kerosene (3 samples)	CONCAWE 91/51
	Straight run kerosene	Blackburn et al '86
	Hydrotreated kerosene	Blackburn et al '86
<b>Mouse lymphoma assay</b>		
	Straight run kerosene	API 32-32745
	Straight run kerosene	API 26-60017
	Hydrodesulfurized kerosene	API 32-30240*
	Jet Fuel A	API 27-30051
	Jet Fuel JP-5	NTP 1986
<b>Sister chromatid exchange assay (CHO cells)</b>		
	Hydrodesulfurized kerosene	API 35-32482*

**Type** : Ames test

## 5. Toxicity

Id Kerosene

Date 9/21/2010

**System of testing** : S. typhimurium TA1535, TA1537, TA1538, TA98 & TA100 and S. cerevisiae D4  
**Test concentration** : 1/8, 1/4, 1/2 & 1/1 of LD50 equivalent to 0.001 - 5.0 µl/plate  
**Cycotoxic concentr.** : LD50 = 5% in DMSO  
**Metabolic activation** : With and without  
**Result** : Negative  
**Year** : 1977  
**GLP** : No data  
**Test substance** : Kerosene

**Method** : The test solvent used was DMSO.  
The standard Ames plate and suspension assays were conducted with and without metabolic activation. Araclor induced rat liver microsomes were used for metabolic activation.  
Cytotoxicity of the test material was determined prior to the assay i.e the LD<sub>50</sub> was determined and was found to be 5%.  
The concentrations used in the mutagenicity assays are shown below. The lowest concentration used was below a concentration that demonstrated any toxic effects.

Test dose	Percent concentration	
	Bacteria	Yeast
1/8 50% survival	0.625	0.625
1/4 50% survival	1.25	1.25
1/2 50% survival	2.5	2.5
50% survival	5.0	5.0

The following positive control materials were used:

Non-activation assay

Substance	Solvent
Ethylmethane sulfonate	Water or saline
Methylnitrosoguanidine	Water or saline
2-Nitrofluorene	DMSO
Quinacrine mustard	Water or saline

Activation assay

2-Anthramine	DMSO
2-Acetylaminofluorene	DMSO
8-Aminoquinoline	DMSO
Dimethylnitrosamine	Water or saline

**Result** : All results from the plate assays and the suspension assays were negative both in the presence and absence of metabolic activation.

**Test substance** : The sample of kerosene was characterized as follows:

API gravity	41.7°
Initial boiling point	350 °F
Final boiling point	519 °F
Reid vapor pressure	1.7 lb
Sulfur	111 ppm
Vapor density	6.5
Paraffins	32% (v/v)
Olefins	1% (v/v)
Naphthenes	49% (v/v)
Aromatics	18% (v/v)
C6 aromatics	-
C7 aromatics	-
C8 aromatics	-
C9+aromatics	18% (v/v)

**Reliability** : (3) invalid

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

The standard Ames assay is an unsuitable test system for insoluble petroleum products.

(3)

**Type** : Ames test  
**System of testing** : S. typhimurium TA98 & TA100  
**Test concentration** : 10 µl/plate  
**Cycotoxic concentr.** : Not stated  
**Metabolic activation** : With  
**Result** : Negative  
**Year** : 1978  
**GLP** : No data  
**Test substance** : Deodorized kerosene

**Reliability** : (3) invalid  
Report contains few details.  
Furthermore, this assay has been shown to be an inappropriate method for insoluble petroleum distillates.

(4)

**Type** : Ames test  
**System of testing** : S. typhimurium TA 1535, TA1537, TA1538, TA98 & TA100  
**Test concentration** : 245 to 41 000 µg/plate  
**Cycotoxic concentr.** : > 40 000 µg/plate in ethyl acetate  
**Metabolic activation** : With and without  
**Result** : Negative  
**Year** : 1979  
**GLP** : No data  
**Test substance** : Jet fuel A

**Reliability** : (3) invalid  
This assay has been shown to be an inappropriate method for insoluble petroleum distillates.

(5)

**Type** : Modified Ames assay  
**System of testing** : S. typhimurium TA98  
**Test concentration** : 50 µl/plate  
**Cycotoxic concentr.** : No data  
**Metabolic activation** : With  
**Result** : Negative  
**Year** : 1991  
**GLP** : No data  
**Test substance** : Kerosene, 3 samples

**Method** : The method described by Blackburn et al (1986) was used to determine the mutagenicity index of three samples of kerosene. This method differs from the standard Ames assay in the following respects:

- A DMSO extract of the test material is used
- The S9 fraction is obtained from Araclor-induced hamster liver
- An eight-fold concentration of S9 is used
- A two-fold concentration of cofactor NADP is used
- The assay is only conducted in S. typhimurium strain TA98

The DMSO extracts were tested over a range of concentrations that permitted the construction of a dose-response curve.

A Mutagenicity index (MI) was determined for each assay. This was the

## 5. Toxicity

Id Kerosene

Date 9/21/2010

tangent to the dose-response curve at zero dose.  
An assay was judged to be positive if the MI was greater than 1.0

**Result** : The mutagenicity indices of the three samples were all zero, thus demonstrating an absence of mutagenic activity. Interestingly, none of the samples was found to contain any measurable 3-7 ring PACs.

**Test substance** : Three samples were tested. All were straight run kerosenes:

Sample	CAS No.	Treatment	Crude source
1	8008-20-6	wet treated	Middle East
2	64742-81-0	hydrotreated	Middle East
3	64742-81-0	hydrotreated	North Sea

**Reliability** : (4) not assignable  
The results are given as a summary in a review. No experimental details are provided.

(32) (37)

**Type** : Modified Ames assay

**System of testing** : S. typhimurium TA98

**Test concentration** : 50 µl/plate

**Cycotoxic concentr.** : No data

**Metabolic activation** : With

**Year** : 1986

**GLP** : No data

**Test substance** : 2 Samples: Hydrotreated kerosene & Straight run kerosene

**Result** : The mutagenicity index reported for each of the samples was

Straight run kerosene (8008-20-6) MI 2.9

Hydrotreated kerosene (64742-47-8) MI 0.0

**Reliability** : (4) not assignable  
Paper is a compilation of results of many petroleum distillates. No actual data are provided.

(32)

**Type** : Mouse lymphoma assay

**System of testing** : Mouse lymphoma L5178Y cell line

**Metabolic activation** : With and without

**Result** : Negative

**Year** : 1984

**GLP** : Yes

**Test substance** : Hydrodesulfurized kerosene Sample API 81-07

**Method** : Non-Activation assay

Cultures of mouse lymphoma cells were exposed to the test material for four hours at doses that were selected during a cytotoxicity study that had been carried out previously. Following exposure, the cells were washed and placed in growth medium for two or three days to allow recovery, growth and expression of the induced TK-/- phenotype. Cell counts were made daily and appropriate dilutions were made to allow optimal growth rates.

At the end of the expression period,  $3 \times 10^6$  cells for each dose were seeded in soft agar plates with selection medium and resistant (mutant) colonies were counted after 10 days incubation. To determine the actual number of cells capable of forming colonies, a portion of the cell suspension was also cloned in normal (non-selective) medium. The ratio

## 5. Toxicity

Id Kerosene

Date 9/21/2010

of resistant colonies to total viable cell number is the mutant frequency.

### Activation Assay

The activation assay was run concurrently with the non-activation assay. The only difference was the addition of the S9 fraction of rat liver homogenate and necessary co factors during the four hour treatment period.

The final concentrations of the activation system components in the cell suspension were:

2.4 mg NADP/ml; 4.5 mg isocitric acid/ml; 50 µl S9/ml.

S9 homogenate was obtained from Araclor-induced rat liver.

### Evaluation criteria

The minimum condition considered necessary to demonstrate mutagenesis for any given treatment is a mutant frequency that exceeds 150% of the concurrent background frequency by at least  $10 \times 10^{-6}$

### Result

: The test material was immiscible with water and DMSO at 100 µl/ml but was miscible with ethanol at the same concentration.

Stocks were prepared by performing serial dilutions in ethanol just prior to each assay. The mutation assays were then initiated by performing final dilutions of the stocks into the assay medium containing the lymphoma cells. The test material appeared soluble in the assay medium up to 125 nl/ml but a white precipitate was noted from 250 to 1000 nl/ml.

Two trials of the assay were initiated.

Trial 1 was performed with and without activation but the non-activation assay was NOT used in the evaluation because of unacceptable suspension growth in the negative controls.

The non-activation portion of the assay was therefore repeated in Trial 2.

The report summarized here included the acceptable activation assay from Trial 1 and the acceptable non-activation assay from Trial 2.

The results are summarized below.

	Rel Susp. growth (% of control)	Total mutant colonies	Total viable eff.	Cloning eff.	Rel growth (%)	Mutant frequency $10E^{-6}$ units
<b>Non activation assay (Trial 2)</b>						
Solvent control (ethanol)						
	100	70	392	100	100	17.9
	100	74	287	100	100	25.8
	100	75	367	100	100	20.4
	100	50	324	100	100	15.4
Untreated control						
	107.1	82	374	109.2	116.7	21.9
	97.4	70	392	116.5	111.3	17.9
EMS (µl/ml)						
0.5	46.7	851	170	49.6	23.2	500.6
0.5	57.3	753	155	45.3	25.9	485.8
API 81-07 (nl/ml)						
6.25	28.3	83	331	96.6	27.3	25.1
12.5	24.7	61	282	82.3	20.3	21.6
12.5	67.4	89	332	96.9	65.3	26.8



## 5. Toxicity

Id Kerosene

Date 9/21/2010

25.0	16.3	87	258	75.3	12.3	33.7
25.0	45.4	61	248	72.4	82.7	24.6
37.5	10.3	105	366	106.8	11	28.7
37.5	6.2	77	143	41.7	2.6	53.8
<b>Activation assay</b>						
Solvent control (ethanol)						
	100	98	236	100	100	41.5
	100	92	282	100	100	32.6
Untreated control						
	128.2	99	226	87.2	111.8	43.8
DMN ( µl/ml)						
	0.3	58.5	151	34	13.1	7.7
						444.1
API 81-07 (nl/ml)						
	3.91	100.1	66	194	74.8	74.9
						34
	7.81	194.1	52	144	55.6	107.9
						36.1
	15.6	101.7	42	259	99.9	101.6
						16.2
	31.3	27.3	45	158	61	16.7
						28.5
	62.5	9	68	175	67.5	6.1
						38.9

Under non-activation conditions the test material induced a good range of toxicities for evaluation (relative growths ranged from 2.8% to 65.3%). None of the assays induced a mutant frequency that exceeded the minimum criterion ( $40.8 \times 10^{-6}$ ). The test material was not mutagenic under non-activation conditions.

In the presence of metabolic activation a wide range of toxicities was induced (6.1 to 107.9% relative growths). The minimum criterion mutant frequency of  $69.0 \times 10^{-6}$  was not exceeded. The test material was therefore considered non mutagenic under activation conditions.

**Reliability** : (1) valid without restriction (14)

**Type** : Mouse lymphoma assay  
**System of testing** : Mouse lymphoma cells L5178Y cell line  
**Test concentration** : 0.5 to 00067 µl/ml  
**Cycotoxic concentr.** : 100% mortality at 0.075 µl/ml, 10% mortality at 0.01 µl/ml  
**Metabolic activation** : With and without  
**Result** : Positive  
**Year** : 1985  
**GLP** : Yes  
**Test substance** : Straight run kerosene, API sample 83-09

**Reliability** : (1) valid without restriction (17)

**Type** : Mouse lymphoma assay  
**System of testing** : Mouse lymphoma cells L5178Y cell line  
**Test concentration** : With activation: 0.004 to 0.065 µl/ml, without activation: 0.006 to 0.13 µl/ml  
**Cycotoxic concentr.** : Toxic at 0.13 µl/ml, slightly toxic at 0.065 µl/ml  
**Metabolic activation** : With and without  
**Result** : Negative  
**Year** : 1977  
**GLP** : No data  
**Test substance** : Straight run kerosene  
**Reliability** : (1) valid without restriction (3)

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

**Type** : Mouse lymphoma assay  
**System of testing** : Mouse lymphoma cells L5178Y cell line  
**Test concentration** : With activation:25 to 200 µg/ml, without activation 100 to 1200 µg/ml  
**Cytotoxic concentr.** : Max concentration that could be used was 0.1%  
**Metabolic activation** : With and without  
**Result** : Positive  
**Year** : 1979  
**GLP** : No data  
**Test substance** : Jet fuel A

**Result** : Negative in the non-activation assay.  
Positive in the activation assay.

**Reliability** : (1) valid without restriction

(5)

**Type** : Mouse lymphoma assay  
**System of testing** : Mouse lymphoma cells L5178Y cell line  
**Test concentration** : 10 mg/plate  
**Result** : negative  
**Year** : 1986  
**Test substance** : Jet fuel JP-5

(61)

**Type** : Sister chromatid exchange assay  
**System of testing** : Chinese Hamster Ovary cells  
**Metabolic activation** : With and without  
**Result** : Negative  
**Year** : 1988  
**GLP** : Yes  
**Test substance** : Hydrodesulfurized kerosene, API sample 81-07

**Method** : A cytotoxicity study was performed in order to select dose levels for the SCE assay.  
For the SCE assay CHO cells were seeded in duplicate for each treatment condition and were incubated at 37°C in a humidified atmosphere for 16 to 24 hours.  
Treatment was carried out by refeeding two complete sets of flasks with complete medium for the non activation study or with S-9 reaction mixture for the activated study to which was added 50 µl of dosing solution of test control or article in solvent or solvent alone.  
In the non activation study the cells were exposed for about 25 hours. At the end of the treatment period, the treatment medium was removed, the cells rinsed and then exposed to 0.01mM BrdUrd and colcemid (0.1 µg/ml) for a further 2 hours.  
In the activation study exposure was for 2 hours.  
After the exposure period, the treatment medium was removed, the cells were washed re-fed with medium containing BrdUrd and then incubated for a further 26 hours. Colcemid was added for the last 2 hours of incubation. For activated and non activated assays metaphase cells were harvested 2 hours after addition of colcemid. Cells were collected and fixed and stored until slides were prepared.  
Slides were coded and scored without regard to treatment group. Only cells with 20 ±2 centromeres were selected for evaluation of SCEs. A total of 4 doses were scored including the highest test article dose where sufficient second-division metaphase cells were available. SCEs were scored in 25 cells from each duplicate culture to make up a total of 50 cells

## 5. Toxicity

Id Kerosene

Date 9/21/2010

per treatment. The percentage of cells in first (M1), second (M2) or third division (M3) metaphase was also recorded for a total of 100 metaphase cells scored.

TEM was used as positive control in the non activated assay at a concentration of 0.025 µg/ml. CP was used in the activation assay at a concentration of 2.5 µg/ml.

S-9 was prepared from Araclor-induced rat liver.

The test material was considered positive if it induced a doubling in SCE frequency over the solvent control at a minimum of three consecutive dose levels or if a dose responsive and statistically significant increase was observed over a minimum of three dose levels.

### Result

: The test material was soluble at all concentrations tested. The study in both the presence and absence of S9 was repeated since there was a poor metaphase cell yield. Only the results of the second study with and without S9 are summarized in the following table.

Replicate flask	SCEs/ chromosome	Flask mean SCEs/cell (±SD)	Group mean SCEs/cell (±SD)
<b>Assay in absence of exogenous activation</b>			
Untreated cells			
A	0.42	8.4±3.16	
B	0.42	8.28±2.57	8.34±2.85
Acetone			
A	0.48	9.44±3	
B	0.45	8.96±2.35	9.20±2.68
API 81-07 0.007 µl/ml			
A	0.44	8.80±2.87	
B	0.43	8.68±2.54	8.74±2.69
API 81-07 0.013 µl/ml			
A	0.47	9.4±2.96	
B	0.42	8.32±2.58	8.86±2.80
API 81-07 0.025 µl/ml			
A	0.47	9.36±2.96	
B	0.48	9.64±3.12	9.50±3.01
API 81-07 0.05 µl/ml			
A	NE <sup>(a)</sup>		
B	NE		
TEM			
A	1.53	30.6±6.81	
B	1.75	34.92±7.60	2.76±7.47**

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

### Assay in presence of exogenous activation

#### Untreated cells

A	0.5	10.16±2.98	
B	0.55	11.04±2.76	10.6±2.88

#### Acetone

A	0.47	9.36±3.94	
B	0.45	8.96±2.99	9.16±3.47

#### API 81-07 0.05 µl/ml

A	0.63	12.52±4.09	
B	0.58	11.64±3.38	12.08±3.74**

#### API 81-07 0.1 µl/ml

A	0.48	9.56±3.8	
B	0.51	10.36±4.27	9.96±4.02

#### API 81-07 0.2 µl/ml

A	0.5	10.04±3.23	
B	0.44	8.84±3.5	9.44±3.39

#### API 81-07 0.4 µl/ml

A	0.5	9.96±3.25	
B	0.58	11.56±3.57	10.76±3.47*

#### CP

A	1.91	38.2±7.06	
B	2.01	40.2±12.18	39.2±9.91**

(a) Not evaluated due to absence of second-division metaphase cells

\*  $P \leq 0.05$

\*\*  $P \leq 0.01$

The responses to the positive and negative control materials fulfilled the requirements for the assays.

The test material did not cause an increase in SCEs in the absence of exogenous activation.

API 81-07 did cause a significant increase in SCEs at two non adjacent doses in the activation assay. However, the increased activity was only seen in one of two treatment flasks. These increases appeared to be random and of no biological significance.

It was concluded that API 81-07 was negative in the SCE assay.

#### Reliability

: (1) valid without restriction

(24)

## 5. Toxicity

Id Kerosene

Date 9/21/2010

**Type** : Overall summary of in-vitro studies  
**System of testing** : Various

**Result** : Summarized results of all the reported in vitro studies of genotoxicity are shown below

Test material	Result		Reference
	+S9	-S9	
Standard Ames assay			
Deodorized kerosene	-	-	API 26-60103
Straight run kerosene	-	-	API 26-60017*
Jet Fuel A	-	-	API 27-30051
Modified Ames assay			
Straight run kerosene (MD5)			
	-	ND	CONCAWE 91/51*
Straight run kerosene	+	ND	Blackburn et al 86*
Hydrotreated kerosene (MD 2, 3, & 4)			
	-	ND	CONCAWE 91/51
Hydrotreated kerosene	-	ND	Blackburn et al '86
Mouse Lymphoma assay			
Straight run kerosene (83-09)			
	± <sup>(a)</sup>	+	API 32-32745
Straight run kerosene	-	-	API 26-60017
Hydrodesulfurized kerosene (81-07)			
	-	-	API 32-30240*
Jet Fuel A	+	-	API 27-30051
Jet Fuel JP-5	-	-	NTP 1986
Sister chromatid exchange assay (CHO cells)			
Hydrodesulfurized kerosene (81-07)			
	-	-	API 35-32482*

<sup>(a)</sup> Equivocal with activation

\* Studies for which full robust summaries have been prepared

### 5.5 GENETIC TOXICITY *IN VITRO*

#### TEST SUBSTANCE

**Category Chemical:** Kerosene/Jet Fuel

**Test Substance:** JP-8 Jet Fuel

**Test Substance Purity/Composition and Other Test Substance Comments:** JP-8 supplied by Wright-Patterson Air Force Base. No other details provided.

**Category Chemical Result Type:**

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

### Unable to Measure or Estimate Justification:

#### METHOD

**Type of Study:** In vitro exposure of H4IIE rat hepatoma cells followed by comet (single cell gel electrophoresis) assay.

**Concentration:**

**Concentrations:** H4IIE rat hepatoma cells were exposed to 3, 5, 10, 15 and 20 mg/ml JP-8 (dissolved in 0.1% EtOH) for 4 h.

**Year Study Performed:** 2000

**Method/Guideline Followed:** Comet assays were performed using alkaline unwinding of DNA and neutral electrophoresis following the protocol for rapid detection of DNA damage from the CometAssay™ kit (Trevigen Inc., Gaithersburg, MD).

**GLP:** Unknown

**Positive, Negative and Solvent Control Substance(s):** Negative (media) control and solvent (0.1% ethanol) control used.

**Method/Guideline and Test Condition Remarks:**

#### TEST RESULTS

**Details for Cytogenetic Assay (if applicable):**

**Statistics:** The Kruskal–Wallis one-way analysis of variance by ranks was used as a non-parametric test to determine whether the distributions of the various tail parameters differed in exposure groups within a fuel treatment. Statistically significant results were subjected to the Dunn's post test ( $\alpha = 0.05$ ) to compare the differences in the groups with the expected average difference.

<b>Species:</b>	<b>Strain:</b>	<b>Metabolic Activation:</b>	<b>Genotoxic Effect:</b>	<b>Conclusion:</b>
Rat	H4IIE rat hepatoma cells	No exogenous source used	Comet assay	Positive

**Other Species**

**Other Strain:**

**Other Species:**

**Other Strain:**

**Other Species:**

**Other Strain:**

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

Other Species:

Other Strain:

Other Species:

Other Strain:

Other Species:

Other Strain:

Results Remarks:

Conclusion Remarks:

Exposure to JP-8 resulted in an overall increase in mean comet tail moments ranging from 0:74 \_ 0:065 (0.1% EtOH control) to 3:13 \_ 0:018; 4:36 \_ 0:32; 5:40 \_ 0:29; 7:70 \_ 0:52 and 11:23 \_ 0:77 for JP-8 concentrations 3, 5, 10, 15 and 20mg/ml, respectively. Addition of DNA repair inhibitors hydroxyurea (HU) and cytosine arabinoside (Ara-C) to cell culture with JP-8 resulted in accumulation of DNA damage strand breaks and increase in comet tail length. Inclusion of 4mM HU and 40mM Ara-C with 3, 5, 10 and 20mg/ml JP-8 concentrations resulted in increased mean tail moments to 5:94 \_ 0:43; 10:12 \_ 0:72; 17:03 \_ 0:96; and 29:25 \_ 1:55. JP-8, in the concentrations used in this study, did not result in cytotoxicity or significant apoptosis, as measured using the terminal deoxynucleotidyl transferase (TDT)-mediated dUTP-X nick end labeling (TUNEL) assay. These results demonstrate that relevant exposures to JP-8 result in DNA damage to H4IIE cells, and suggest that DNA repair is involved in mitigating these effects.

### RELIABILITY/DATA QUALITY

Reliability: 2

Reliability Remarks: Reliable with restrictions

Key Study Sponsor Indicator:

### REFERENCE

Reference:

Geraldine M. Grant, Shawna M. Jackman, Christopher J. Kolanko, David A. Stenger. JP-8 jet fuel induced DNA damage in H4IIE rat hepatoma cells. Mutation Research 490 (2001) 67–75

## 5.5 GENETIC TOXICITY *IN VITRO*

### TEST SUBSTANCE

Category Chemical: Kerosene/Jet Fuel

Test Substance: Distillates (petroleum), hydrotreated light (CAS #64742-47-8)

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

Test substance met the specifications for commercial Jet A (aviation turbine fuel, ASTM D 1655).

**Test Substance Purity/Composition and Other Test Substance Comments:**

Test sample	Boiling range	Aromatic content
4+rings aromatics (wt. %)		
Turbo fuel A	326- 524°F (163-273°C)	21.8 vol %
ND		

**Category Chemical Result Type:**

**Unable to Measure or Estimate Justification:**

### METHOD

**Type of Study:**

In vitro *Salmonella typhimurium* mutagenesis test

**Concentration:**

The aromatic fractions of the test sample was prepared by having one ml of test materials extracted directly with 5 ml DMSO (the DMSO extract). It should be noted that at no point in the extraction procedure is the DMSO-extractable material concentrated and weighed, so it is only possible to relate the material tested to the starting material in very general terms. However, in the extraction process, there is a 5-to-1 ratio of DMSO to starting material. Thus each uL DMSO extract contains the extractable constituents from 0.2 ul of the starting material. The density of middle distillate fuels is approximately 0.8 mg/cc. Thus the 1-50 uL of material tested in the cyclohexane extract study represented the extractable material from -0.16 – 8.0 mg of the starting material, and in the DMSO extract study, the 5-60 uL of material tested represented approximately 0.8-9.6 mg.

**Concentrations:**

**DMSO Extract Dose (uL/plate)**  
60 (toxic)  
50 (toxic)  
40  
30  
20  
15  
10  
5

**Year Study Performed:**

1994

**Method/Guideline Followed:**

A modification of the standard *Salmonella typhimurium* mutagenesis test [Ames et al., 1975] was developed by Blackburn and coworkers (1984,1986). Three changes were made in the standard assay. The first was the use of an extraction step to concentrate the potentially mutagenic aromatic components. The second was the use of S-9 from Aroclor-induced hamster liver at 8 times the recommended concentration. The third was the exclusive use of TA98, the tester strain most responsive to complex mixtures of PAH. The testing involved a preincubation procedure. The components were added in the following order: 60 ul DMSO extract, 400 ul S-9 mixture.



# 5. Toxicity

**Id** Kerosene  
**Date** 9/21/2010

And 0.1 ml *Salmonella*. Test material aliquots contained between 1 and 60 ul of extract in a total volume of 60 ul DMSO was used as the diluent. The mixture was then incubated for 20 min at 37°C. Following the incubation period, top agar was added, and the mixture was then poured onto petri plates. After a 72-hr incubation period at 37°C. the plates were scored for revertants.

**GLP:** Unknown

**Positive, Negative and Solvent Control Substance(s):** Positive control was 2-aminoanthracene and the negative control was DMSO

**Method/Guideline and Test Condition Remarks:** Method used was later adopted by ASTM as E 1687

## TEST RESULTS

**Details for Cytogenetic Assay (if applicable):**

Each dilution of test material was plated in triplicate. If revertant levels at all doses were less than twice control values, the sample was judged to be non-mutagenic. For samples producing revertant frequencies more than twice background levels, the initial slope of the dose-response curve (i.e., the mutagenic index, MI) was determined by the method of Myers [1981].

**Statistics:**

Species:	Strain:	Metabolic Activation:	Genotoxic Effect:	Conclusion:
<i>Salmonella typhimurium</i>	T98	Yes. Aroclor induced hamster S9	Mutation	Negative
<b>Other Species:</b>				
<b>Other Strain:</b>				
<b>Other Species:</b>				
<b>Other Strain:</b>				
<b>Other Species:</b>				
<b>Other Strain:</b>				
<b>Other Species:</b>				
<b>Other Strain:</b>				
<b>Other Species:</b>				

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

Other Strain:	
Other Species:	
Other Strain:	

### Results Remarks:

**Conclusion Remarks:** Turbo fuel A produced less than a 2-fold increase in revertant colonies when compared to the appropriate vehicle controls. Thus under the conditions of this test, these materials were not considered mutagenic, and mutagenic indices were not calculated.

### RELIABILITY/DATA QUALITY

**Reliability:** 2

**Reliability Remarks:** Reliable with restrictions

**Key Study Sponsor Indicator:**

### REFERENCE

**Reference:** R.H. McKee, M.A. Amoruso, J.J. Freeman, and R.T. Przygoda. Evaluation of the Genetic Toxicity of Middle Distillate Fuels. Environmental and Molecular Mutagenesis 23:234-238 (1994)

## 5.6 GENETIC TOXICITY 'IN VIVO'

**Type** : In vivo studies

**Remark** : The in vivo genotoxic potential of kerosenes and jet fuel has been studied in several studies and these are tabulated below. Only one study for each test system is described in detail in the following section on in vivo genotoxicity studies and these are indicated (\*) in the table. For completeness, the results of all studies are also briefly summarized as separate robust summaries. At the end of this section the results of all in vivo studies are summarized in tabular form.

<u>Assay</u>	<u>Reference</u>
<b>Rat bone marrow cytogenetics</b>	
Hydrodesulfurized kerosene (81-07)	API 32-30240*
Straight run kerosene (83-09)	API 32-31769
Straight run kerosene (8008-20-6)	API 26-60017
Straight run kerosene (8008-20-6)	API 26-60017
Jet fuel A	API 27-30051*
<b>Sister chromatid exchange</b>	
Hydrodesulfurized kerosene (81-07)	API 36-30043*
<b>Dominant lethal assay</b>	
Deodorized kerosene	API 26-60098
Jet fuel A	API 28-31345*

\* Full robust summaries prepared for these studies

**Type** : Cytogenetic assay  
**Species** : Rat  
**Sex** : Male/female  
**Strain** : Sprague-Dawley  
**Route of admin.** : i.p.  
**Doses** : 0.3, 1.0 & 3.0 g/kg  
**Result** : Negative  
**Year** : 1984  
**GLP** : Yes  
**Test substance** : Hydrodesulfurized kerosene Sample API 81-07

**Method** : A pilot study was carried out in 4 male and 4 female young adult Sprague Dawley rats. These animals were given a single intraperitoneal (i.p.) dose (3 g/kg) of API 81-07. During the following 48 hours observation, no animals died. The doses selected for the cytogenetics study were therefore 0.3, 1 and 3 g/kg. Three groups of 15 male and 15 female rats were given a single i.p. dose of either 0.3, 1 or 3 g API 81-07/kg. At six, 24 and 48 hours after dosing 5 males and 5 females were killed at each dose level. An additional 15 males and 15 females were untreated and served as negative controls. These animals were otherwise treated the same as the test animals. A positive control group of 5 males and 5 females was administered 0.8 mg/kg Triethylenemelamine (TEM) as a single i.p. dose. These positive control animals were killed 24 hours after administration of the positive

control substance.

Three hours prior to being killed with CO<sub>2</sub>, animals were injected i.p. with 4 mg/kg of colchicine. After the animal was killed, the adhering soft tissue and epiphyses of both tibiae were removed and the marrow was flushed from the bone and transferred to Hank's balanced salt solution. The marrow button was collected by centrifugation and was then re suspended in 0.075M KCl. The centrifugation was repeated and the pellet re suspended in fixative (methanol:acetic acid, 3:1). The fixative was changed once and left overnight. Cells in fixative were dropped onto glass slides which were then air dried and stained with Giemsa. Slides were coded and scored for chromosomal aberrations.

50 spreads were read for each animal where feasible.

A mitotic index based on at least 500 counted cells was also recorded. The index was calculated by scoring the number of cells in mitosis per 500 cells on each read slide.

Statistical evaluation

Performed by Student's t-tests on four parameters:

1. Number of structural aberrations per animal
2. Number of numerical aberrations per animal
3. % cells with one or more structural aberrations per animal
4. % cells with 2 or more structural aberrations per animal.

Data interpretation and evaluation

Gaps were not counted as significant aberrations.

Open breaks were considered as indicators of genetic damage as were configurations resulting from the repair of breaks. The latter included translocations, multiradials, rings, multicentrics, etc. Reunion figures such as these were weighed slightly higher than breaks since they usually resulted from more than one break.

Cells with more than one aberration were considered to indicate more genetic damage than those with evidence of single events.

Consistent variations from the euploid number were also considered in the evaluation of mutagenic potential.

The type of aberration, its frequency and its correlation to dose in a given time was considered in evaluating the test material as being positive or negative.

**Result** : The data are given in the report for males, females and as male and female pooled data.

When the results for males were compared with those for controls and the females were compared to controls, no statistically significant differences were found. The data summarized below, are the pooled data for males and females.

Time after dose (hrs)	No of rats	Total No of cells	% cells with aberrations		Mitotic index
			1+	2+	
Negative control					
6	6	300	0	0	5.6
24	10	500	0.4	0	4.4
48	10	500	0	0	4.3
Positive control (TEM, 0.8 mg/kg)					
24	7	253	19.4**	10.7*	1.0

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

API 81-07, 0.3 g/kg  
6 8 292 0.7 0 4.2  
24 9 450 0 0 4.5  
48 10 500 0 0 4.9

API 81-07, 1.0 g/kg  
6 8 400 0.8 0.2 5.2  
24 9 450 0 0 4.4  
48 9 407 1.0 0.2 7.0

API 81-07, 3.0 g/kg  
6 8 365 0.3 0 4.0  
24 10 475 0.6 0 3.5  
48 8 400 0 0 6.0

\* Significant if outlier excluded  
\*\*P < 0.01

The structural aberration frequency did not differ significantly from the negative control at any tested dose. The percentage of cells showing one or more structural aberrations or 2 or more structural aberrations were also similar to the negative controls. A concurrent positive control group induced significant increases in aberrations.

**Reliability** : (1) valid without restriction (14)

**Type** : Cytogenetic assay  
**Species** : Rat  
**Sex** : Male/female  
**Strain** : Sprague-Dawley  
**Route of admin.** : i.p.  
**Exposure period** : 6, 24 and 48 hours  
**Doses** : 0, 0.3, 1.0 & 3.0 g/kg  
**Result** : Negative  
**Year** : 1985  
**GLP** : Yes  
**Test substance** : Straight run kerosene, API sample 83-09

**Reliability** : (1) valid without restriction (18)

**Type** : Cytogenetic assay  
**Species** : Rat  
**Sex** : Male/female  
**Strain** : Sprague-Dawley  
**Route of admin.** : i.p.  
**Exposure period** : 6, 24 & 48 hours  
**Doses** : Sample 1: 0.4, 0.13 & 0.04 ml/rat; Sample 2: 0.18, 0.06, 0.02 ml/rat  
**Result** : Negative  
**Year** : 1977  
**GLP** : Yes  
**Test substance** : Straight run kerosene, 2 samples

**Result** : Neither of the samples cause aberrations in bone marrow cells.  
**Reliability** : (1) valid without restriction (3)

## 5. Toxicity

Id Kerosene

Date 9/21/2010

**Type** : Sister chromatid exchange assay  
**Species** : Mouse  
**Sex** : Male/female  
**Strain** : B6C3F1  
**Route of admin.** : i.p.  
**Exposure period** : 20-22 hours  
**Doses** : 400, 2000 & 4000 mg/kg  
**Result** : Positive  
**Year** : 1988  
**GLP** : Yes  
**Test substance** : Hydrodesulfurized kerosene, sample API 81-07

**Method** : Six groups of five male and five female, 8-10 week old B6C3F1 mice were anesthetized with Metofane and an agar-coated BrdUrd pellet was implanted subcutaneously in the lower abdominal region.

Four hours after implantation of the pellet the mice were given test material as a single i.p dose at doses of either 400, 2000 or 4000 mg/kg in corn oil at a dose volume of 10 ml/kg (doses based a range finding study). A solvent control group was given corn oil at a dose of 10 ml/kg. Two positive control groups were used. One received cyclophosphamide i.p. at a dose of 10 ml/kg and the other received API 81-15 (a carcinogenic heavy fuel oil component) at a dose level of 4 g/kg, also at an injection rate of 10 ml/kg. Two to four hours prior to sacrifice, colchicine was injected intraperitoneally at a dose of 1 mg/kg to arrest dividing cells at metaphase. 24-26 hours after pellet implantation all mice were sacrificed by CO<sub>2</sub>. Both femurs of each animal were cut and the marrow was aspirated into a syringe containing cold Hank's balanced salt solution. The cells were collected by centrifugation, resuspended in 5 ml warm hypotonic solution and then incubated for approximately 10 minutes at 37 °C to swell the cells. The swollen cells were collected by centrifugation, resuspended in two consecutive changes of methanol/acetic acid (fixative) and then stored in a refrigerator prior to slide preparation. Slides were prepared by resuspending the cells in fresh fixative, by centrifugation and then dropping 2 to 4 drops of fixed cells onto a wet slide. The slides were then air-dried. Two to 5 slides were prepared from each animal. The slides were stained with Giemsa for subsequent examination.

A minimum of 50 second-division metaphase spreads from each animal were examined and scored for SCEs and chromosome number. The mitotic index was recorded as the % of cells in mitosis based upon 500 counted cells. The % first, second and third-division metaphase cells was also recorded as the number per 100 counted cells.

Using the number of animals per group as the sample size, statistical analysis was performed with the Kruskal Wallis test and, if required, the Mann Whitney test.

The test article was considered to induce a positive response if a dose-related increase in SCEs/metaphase was observed relative to the vehicle control (P<0.05, Mann Whitney test).

Criteria for determination of a valid test

The mean number of SCE per second-division metaphase cell must not exceed 8 SCEs/cell in the negative (vehicle) control. The mean SCE/metaphase for the positive control animals must be statistically

## 5. Toxicity

Id Kerosene

Date 9/21/2010

### Result

- increased relative to the vehicle control using the Mann Whitney test (P<0.05).
- : The high dose male mice were lethargic shortly after dose administration and on the following day. In the high dose males and females and the mid-dose males there was a slight weight loss between the time pretreatment body weights were measured and when the animals were treated with colchicine on the following day.
- The body weight changes are shown below:

Treatment	Sex	% change (+ or -) 24 hr
Corn oil 10 ml/kg	M	0
	F	- 2.9
API 81-07 400 mg/kg	M	- 1.4
	F	- 0.5
2000 mg/kg	M	- 3.5
	F	- 1.9
4000 mg/kg	M	- 4.3
	F	- 4.7
API 81-15 4000 mg/kg	M	0
	F	+ 3.8
Cyclophosphamide 10 mg/kg	M	+ 2.6
	F	- 3.8

The SCEs were counted in 50 second-division metaphase cells and these data are summarized in the following table.

Treatment	Sex	Range of mean SCEs/cell for individual animals	Average SCEs/cell per mouse <sup>(a)</sup>
Corn oil	M	4.68 - 5.84	5.44±0.47 (5.64)
	F	5.28 - 7.36	6.25±0.86 (6.06)
API 81-07 400 mg/kg	M	6.76 - 10.46	8.26±1.51 (5.64)**
	F	6.98 - 8.26	7.56±0.52 (7.56)
2000 mg/kg	M	5.68 - 7.58	6.74±0.87 (6.86)**
	F	6.8 - 9.84	8.26±1.22 (8.54)
4000 mg/kg	M	5.72 - 7.66	6.86±0.75 (7.06)*
	F	6.56 - 12.3	9.20±2.49 (9.88)
API 81-15 4000 mg/kg	M	6.68 - 9.28	7.94±0.93 (7.94)**
	F	7.28 - 8.54	7.86±0.58 (7.56)*
Cyclophosphamide 10 mg/kg	M	36.6 - 44.18	40.3±3.53 (38.5)**
	F	18.34 - 31.64	25.5±5.44 (25.06)**

## 5. Toxicity

Id Kerosene

Date 9/21/2010

(a) Mean  $\pm$  standard deviation (Median SCEs/cell)  
\* P < 0.05  
\*\* P < 0.01 (Mann Whitney test)

**Reliability** : (1) valid without restriction (23)

**Type** : Dominant lethal assay  
**Species** : Mouse  
**Sex** : Male  
**Strain** : CD-1  
**Route of admin.** : Inhalation  
**Exposure period** : 6 hours each day, 5 days/week for 8 weeks  
**Doses** : Nominal: 0, 100 & 400 ppm. Actual: 0, 98.4 & 378.3 ppm  
**Result** : Negative  
**Method** :  
**Year** : 1980  
**GLP** : No data  
**Test substance** : Jet Fuel A

**Method** : Groups of twelve male mice (14 weeks old) were exposed by inhalation to nominal vapor concentrations of 100 or 400 ppm of test material. Exposures were for 6 hours a day, five days each week for eight weeks. A control group of mice were placed in the exposure chamber each day and were exposed only to filtered air. On day 40 of the dosing schedule, the animals which served as positive controls received a single i.p. injection of 0.3 mg/kg triethylenemelamine (TEM). The mice were observed for clinical signs of toxicity twice daily during the exposure phase of the study. The males were then mated with two females for one week. After mating, the females were removed and housed in other cages. After 2 days rest the males were mated with a further two females for one week. Two weeks after mating, the females were killed using CO<sub>2</sub> and were necropsied. During necropsy, the uteri of each pregnant female were examined and the number of living and dead implants were counted.

The following parameters were calculated:  
Fertility index  
Computed as No. pregnant females per No. mated females.  
Ratio was evaluated by a Chi square test to compare each treatment group and the positive control to the negative control. Armitage's trend for linearity was used to test whether the fertility index was linearly related to arithmetic or log dose.

### Total number of implants

This number was evaluated by Student's t-test to determine whether the average number of implantations per pregnant female for each treatment group and positive controls were different to the negative controls.

### Dead implants

The dead implant frequencies were transformed (Freeman-Turkey), and the transformed data was compared to the negative control group by Student's t-test

Proportion of females with one or more dead implantations  
Proportion of females with two or more dead implantations



## 5. Toxicity

Id Kerosene  
Date 9/21/2010

These two parameters were evaluated similarly. The quotient was evaluated by the same statistical method used for deriving fertility index with the addition of a probit regression analysis to determine if the probit of the proportions is related to log dose.

### Dead implants/Total implants

These were computed for each female and transformed using Freeman-Turkey arc-sine transformation prior to being evaluated by Student's t-test.

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

**Result** : There were no deaths during the exposure period, nor were there any Jet-fuel-related clinical signs. Growth rates were unaffected by exposure. The results obtained from the females are as follows.

	Negative control		Positive control		100 ppm		400 ppm	
	wk1	wk2	wk1	wk2	wk1	wk2	wk1	wk2
Fertility index	.086	.682	.675	.417	.909	.762	.818	.682
Av. No. implantations/pregnant female	11.053	12.533	9.048**	8**	12.5	12.375	12	12.6
Average resorptions (dead implants/pregnant female)	.632	.533	4.619**	5.6**	.95	.875	.222	.4
Proportion of females with one or more dead implantations	.579	.4	.905**	1.0**	.5	.5	.111	.333
Proportion of females with two or more dead implantations	.063	.133	.857**	1**	.4*	.125	.056	.067

\* P < 0.05  
\*\* P < 0.01

With the exception of the proportion of females with two or more dead implantations there were no differences in the developmental parameters in either the one week or two week animals that could be attributable to Jet Fuel A. The one significant finding in the animals mated for one week was considered by the authors to be spurious.

The positive control consistently gave significant increases in the various measured/calculated parameters .

The results support the conclusion that Jet fuel A did not increase the incidence of post-implantation deaths.

**Test condition** : Jet Fuel A vapor was generated by bubbling dry, oil-free, breathing quality air through a column of liquid Jet fuel A in a 500 ml dust impinger in a heated water bath. The concentrated vapor was diluted with room air in the chamber. The concentration in the chamber was determined at least hourly

The nominal and actual concentrations were:

Concentration (ppm)	
Nominal	Actual
0	0
100	98.4
400	378.3

**Reliability** : (1) valid without restriction  
3

(9)

**Type** : Dominant lethal assay  
**Species** : Mouse and rat  
**Sex** : Male  
**Route of admin.** : i.p.  
**Doses** : Mouse: 1 ml/kg (diluted 10% in corn oil), rat: Undiluted  
**Result** : Negative  
**Year** : 1973

## 5. Toxicity

Id Kerosene

Date 9/21/2010

**Test substance** : Deodorized kerosene

(2)

**Type** : Overall summary of in-vivo studies

**Remark** : Summarized results of all the reported in vivo studies of genotoxicity are shown below

<b>Exposure route</b>	<b>Dosage</b>	<b>Results</b>	<b>Reference</b>
<b>Rat bone marrow cytogenetics</b>			
Hydrodesulfurized kerosene (81-07)			
i.p.	0.3, 1.0 & 3 g/kg	negative	API 32-30240*
Straight run kerosene (83-09)			
i.p.	0.3, 1.0 & 3 g/kg	negative	API 32-31769
Straight run kerosene (8008-20-6)			
i.p.	0.02, 0.06 & 0.18 ml/rat 5 consec days	negative	API 26-60017
Straight run kerosene (8008-20-6)			
i.p.	0.04, 0.13 & 0.4 ml/rat	negative	API 26-60017
Jet fuel A			
inhalation	100 ppm for 20 days 400 ppm for 5 days	positive	API 27-30051*
<b>Sister chromatid exchange</b>			
Hydrodesulfurized kerosene (81-07)			
i.p. mouse	0.4, 2.0 & 4.0 g/kg	positive males only	API 36-30043*
<b>Dominant lethal assay</b>			
Deodorized kerosene			
i.p. Mouse & rat	??	negative	API 26-60098
Jet fuel A			
inhalation Mouse	100 & 400 ppm. 6h/day 8 weeks	negative	API 28-31345*

\* Full robust summaries prepared for these studies

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

### 5.6 GENETIC TOXICITY *IN VIVO*

#### TEST SUBSTANCE

**Category Chemical:** Kerosene/Jet Fuel

**Test Substance:** Distillates (petroleum), hydrotreated light (CAS #64742-47-8)

Test substance met the specifications for commercial Jet A turbo fuel (aviation turbine fuel, ASTM D 1655).

#### Test Substance Purity/Composition and Other Test Substance Comments:

Test sample aromatics (wt. %)	Boiling range	Aromatic content	4+rings
Turbo fuel A	326- 524°F (163-273°C)	21.8 vol %	ND

#### Category Chemical Result Type:

#### METHOD

**Type of Study:** In vivo genetic toxicity

**Type of Test:** Mouse bone marrow micronucleus test.

**Route of Administration:** Oral gavage

**Species:** Mouse

**Strain:** CD-1

**Gender:** Male and female

**Dose:** 0.0, 1.0, 2.5, or 5.0 g/kg of test material.

**Year Study Performed:** 1994

**Method/Guideline Followed:** Generally follows OECD 474

**GLP:** Unknown

#### Duration of Treatment/Expososure Period and Units:

**Frequency of Treatment:** One oral dose

**Positive, Negative and Solvent Control Substance(s):** Vehicle control (corn oil) and a positive control (40 mg/kg cyclophosphamide by intraperitoneal injection. All of these mice were sacrificed 24 hr after cyclophosphamide administration).

**Post-Exposure Period:** Five male and five female mice from each group were sacrificed 24, 48, or 72 hr after treatment.

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

**Number of Animals per Sex per Dose:** Five male and five female mice from each group were sacrificed 24, 48, or 72 hr after treatment.

**Method/Guideline and Test Condition Remarks:**

### TEST RESULTS

#### Systemic Toxicity:

Results of CD-1 Mouse Bone Marrow Micronucleus Assays 24-hr Harvest – Males(1)

	Mean PCE (%)	Mean MNE/1,000 PCE
5 g/kg Turbo fuel A	58 ± 4.4	3.2 ± 2.0
2.5 g/kg Turbo fuel A	48 ± 7.9*	1.6 ± 1.5
1 .0 g/kg Turbo fuel A	54 ± 5.2	2.0 ± 1.6
0.04 g/kg Cyclophosphamide	60 ± 2.7	18.6 ± 10.4**
Corn oil (carrier)	56 ± 5.0	1.6 ± 0.9

#### Genotoxic Effect:

1. Similar results were found at 48 hr and 72 hr harvest times, and in female mice at all harvest times.

\*Significantly different from carrier at  $P < 0.05$ )

\*\*Significantly different from carrier at  $P < 0.01$ )

**Results Remarks:** Negative

**Conclusion:** Distillates (petroleum), hydrotreated light (Jet A) was inactive in the mouse micronucleus assay.

### RELIABILITY/DATA QUALITY

**Reliability:** 2

**Reliability Remarks:** Reliable with restrictions

**Key Study Sponsor Indicator:**

### REFERENCE

**Reference:** R.H. McKee, M.A. Amoruso, J.J. Freeman, and R.T. Przygoda. Evaluation of the Genetic Toxicity of Middle Distillate Fuels. Environmental and Molecular Mutagenesis 23:234-238 (1994)

## 5.6 GENETIC TOXICITY *IN VIVO*

### TEST SUBSTANCE

**Category Chemical:** Kerosene/Jet Fuel

**Test Substance:** JP-8 and Jet A (Aviation Turbine Fuels)

**Test Substance Purity/Composition and Other Test Substance Comments:** JP-8 (Lot# 3509) and Jet-A (Lot# 3404) were supplied by the Operational Toxicology Branch, Air Force Research Laboratory, Wright Patterson Air Force Base, Dayton, OH.

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

### Category Chemical Result Type:

#### METHOD

<b>Type of Study:</b>	In vivo genetic toxicity
<b>Type of Test:</b>	Micronucleus studies in the peripheral blood and bone marrow.
<b>Route of Administration:</b>	Dermal
<b>Species:</b>	Mice
<b>Strain:</b>	C3H/HeNCr
<b>Gender:</b>	Female mice, 8–10 weeks old
<b>Dose:</b>	50 – 300 microliters (ul)
<b>Year Study Performed:</b>	2006
<b>Method/Guideline Followed:</b>	Similar to OECD 474
<b>GLP:</b>	Unknown
<b>Duration of Treatment/Exposure Period and Units:</b>	Up to three weeks
<b>Frequency of Treatment:</b>	1 or 3 times, during one or three weeks
<b>Positive, Negative and Solvent Control Substance(s):</b>	Negative (shaved) and positive (cyclophosphamide) controls
<b>Post-Exposure Period:</b>	24 hours
<b>Number of Animals per Sex per Dose:</b>	10
<b>Method/Guideline and Test Condition Remarks:</b>	Two independent investigators (Uni. Texas Health Sciences Center and US Environmental Protection Agency) assessed the incidence of MN in the blood and bone marrow cells. Each investigator used a fluorescence microscope fitted with appropriate filters for acridine orange stain. The incidence of MN was determined from the examination of 2000 consecutive PCE in blood and bone marrow smears (UTHSC and USEPA). The results were decoded after complete microscopic evaluations.
<b>TEST RESULTS</b>	
<b>Systemic Toxicity:</b>	None reported
<b>Genotoxic Effect:</b>	Incidence of MN in PCEs in the blood and bone marrow of mice exposed to JP-8 or Jet-A
	Group      Blood MN/2000 PCEs      Bone marrow MN/2000 PCEs

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

JP-8		
Control	4.4 ± 1.8 (3.1–5.7)	5.1 ± 1.5 (4.0–6.2)
3× 50 ul	4.8 ± 1.4 (3.8–5.8)	4.2 ± 1.8 (2.9–5.5)
3× 100 ul	4.6 ± 1.1 (3.8–5.4)	4.6 ± 0.7 (4.1–5.0)
3× 300 ul	5.0 ± 1.3 (4.1–5.9)	4.8 ± 0.8 (4.2–5.4)
CP	28 ± 4.8*** (24.5–31.5)	25.2 ± 8.2*** (19.3–31.0)
Jet-A		
Control	4.3 ± 1.5 (3.2–5.3)	4.5 ± 1.3 (3.5–5.4)
3× 50ul	5.4 ± 1.8 (4.1–6.7)	3.9 ± 2.9 (1.9–5.9)
3× 100 ul	4.6 ± 1.1 (3.8–5.3)	4.2 ± 0.7 (3.7–4.7)
3× 300 ul	4.8 ± 1.7 (3.6–6.0)	3.9 ± 1.2 (3.0–4.8)
CP	26.4 ± 3.0*** (24.2–28.6)	22.9 ± 7.1*** (17.8–27.9)

Mice were treated dermally with JP-8 or Jet-A on three consecutive days. Blood and bone marrow smears were prepared at 24 h after the last treatment. Similar results were observed with all other dosing regiments. Data are displayed as group mean±S.D and C.I.: confidence intervals at 95% level of the means.

\*\*\* Significant difference between CP (cyclophosphamide) and all other groups,  $p < 0.0001$ .

### Results Remarks:

Whether jet fuel (Jet A or JP-8) was applied dermally once, on three consecutive days, or on a weekly basis showed no statistically significant increases in the incidence of MN/2000 PCE compared to those observed in concurrent untreated control animals. As expected, in all three experiments, significantly elevated frequencies of MN/2000 PCE were observed in positive control (CP-injected) animals.

### Conclusion:

Using several different exposure regimens, no statistically significant increase in the incidence of MN was observed in the bone marrow and/or peripheral blood of mice treated with JP-8 or Jet-A when compared with those of untreated control animals. The data in mice treated with a single dose of JP-8 or Jet-A did not confirm the small but statistically significant increase in micronuclei reported in our previous study[Vijayalaxmi, A.D. Kligerman, T.J. Prihoda, S.E. Ullrich, Cytogenetic studies in mice treated with the jet fuels, Jet-A and JP-8, Cytogenet. Genome Res. 104 (2004) 371–375.]

### RELIABILITY/DATA QUALITY

Reliability: 2

Reliability Remarks: Reliable with restrictions

Key Study Sponsor Indicator:

### REFERENCE

Reference: Vijayalaxmi, A.D. Kligerman, T.J. Prihoda, S.E. Ullrich. Micronucleus studies in the peripheral blood and bone marrow of mice treated with jet fuels, JP-8 and Jet-A. Mutation Research 608 (2006) 82–87.

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

### 5.7 CARCINOGENICITY

**Species** : Mouse  
**Sex** : Male  
**Strain** : C3H  
**Route of admin.** : Dermal  
**Exposure period** : 104 weeks  
**Frequency of treatm.** : 2, 4 or 7 days/week  
**Doses** : Variable, see methods  
**Control group** : Yes, concurrent vehicle  
**Year** : 1996  
**GLP** : Yes  
**Test substance** : Straight run kerosene

**Method** : The testing of the straight run kerosene (MD-3) was part of an overall larger study. For the purpose of this summary, only the details relating to MD-3 are presented.  
The test material was applied to the shorn skin of three groups of 50 male mice for 104 weeks.  
The concentration and dosing frequencies were adjusted to ensure that each animal received the same total weekly dose of test material irrespective of dosing frequency. The concentrations and frequencies were selected to determine the influence of skin irritation on the tumorigenic response.

The following dosing regimes were used

<b>Group No.</b>	<b>Concentration of kerosene* (%)</b>	<b>Dosing frequency <math>\mu</math>l/dose</b>
9	100	50 2 times/week
10	50	50 4 times/week
11	28.5	50 7 times/week

\* Mineral oil was used as diluent

A control group of 50 male mice received 35  $\mu$ l mineral oil 7 days each week. All animals were observed regularly for viability, clinical signs and a score was given for any dermal irritation that occurred. Body weights were recorded throughout the study.

When they developed, dermal growths were measured and documented. All animals were necropsied either when they died during the study or at the end of the study. The necropsy included an examination of the body, all orifices and the carcass, cranial, thoracic and abdominal cavities, including their contents. For all animals, tissues were preserved and examined microscopically from all skin tumors, skin from treated and untreated sites and any grossly observable masses.

**Remark** : The study was designed to assess the carcinogenic potential of middle distillates under conditions in which irritation was present or absent.

The results demonstrated that undiluted kerosene applied twice weekly to the skin caused irritation and skin tumors developed in 12 of 50 mice. In the groups where no irritation occurred, no tumors developed either even though the same total weekly dose of test material was applied to the skin as had been the case with the undiluted material.

**Result** : Survival was less in the 100 % MD-3 treated group compared to the negative controls.  
Dermal irritation occurred in the groups exposed to kerosene.



## 5. Toxicity

Id Kerosene

Date 9/21/2010

The dermal irritation scores were:

Group	Range of scores	Mean dermal score
Negative control (oil)	0-0.22	0.06
100% kerosene 2X/week	0-4.0	2.92
50% kerosene 4X/week	0-0.8	0.09
28.5% kerosene 7X/week	0-0.2	0.04
Positive control (HCO)	0-2.0	0.73

There were no other treatment-related clinical findings.

Treatment did not have any adverse effect on body weights.

Treatment-related findings at post mortem were limited to dermal irritation and were consistent with the findings of the in-life phase of the study.

Liver masses and gastrointestinal abnormalities were observed but these were found in all groups and were considered to be incidental.

Tumors developed in the positive control group (HCO) and only in the MD-3 group that received undiluted sample twice weekly as follows:

Material	Oil	HCO	MD-3	MD-3	MD-3
Concentration			28.5%	50%	100%
No. applications/week			7	4	2
No. mice examined	50	50	50	50	50
No. mice with tumors	0	47	0	0	12
Tumor types					
Squamous cell carcinoma	0	42/73*			7
Spindle cell carcinoma	0	0	0	0	1/2*
Fibrosarcoma	0	0	0	0	3/5*
Melanoma	0	0	0	0	0
Papilloma	0	37/88*	0	0	6

\* / = No with neoplasms/actual incidence of neoplasms

### Test substance

: The sample of straight run kerosene was designated sample MD-3 and had been characterized as follows (CONCAWE 91/51):

CAS No.	64742-81-0
Crude source	Middle East
Kinematic viscosity (cSt)	1.26
Flash point (°C)	49 fire
Distillation range (°C)	175-237 (10 and 90%)
Sulfur content (%)	0.14
Nitrogen content (mg/l)	1.3
Relative density	0.7978
Aniline point	61.4
Aromatics (%)	17.1
Olefins (%)	1.7
Saturates (%)	81.2
PAC 3-7 rings (% wt)	0

### Reliability

: (1) valid without restriction

(37) (47)

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

**Species** : Mouse  
**Route of admin.** : Dermal

**Remark** : The results of several carcinogenicity studies have been reported for kerosene streams and jet fuels. These data have been reviewed and summarized elsewhere (CONCAWE, 1995; IARC, 1989; ASTDR, 1998). The study described in the summary above is considered to be the most recent and the most robust study. However, for completeness, other studies that have been reported are summarized in the following table (taken from CONCAWE, 1995).

Dosing regime	Duration	Result* Tumors	Latency (weeks)	Reference
Straight-run kerosene (API 83-09)				
50 µl, 2x/week 50 mice	>2 years	19 malignant 1 benign	76	API 36-33220
Hydrodesulfurized kerosene (API 81-07)				
50 µl, 2x/week 49 mice	> 2 years	26 malignant 1 benign	77	API 36-31364
Straight run middle distillate (from naphthenic crude)				
50 mg, 2x/week 50 mice	102 weeks	14/30	70	API 32-30964
Straight run middle distillate from paraffinic crude				
50 mg, 2x/week 50 mice	82 weeks	13/27	62	API 32-30964
Jet fuel A				
25 mg, 3x/week 50 mice	105 weeks	11/43	79	Clark 1988
Jet fuel JP-5				
500mg/kg 5x/week 100 mice	103 weeks	No tumors		NTP 1986

It should be noted that in those studies in which tumors developed, moderate to severe skin irritation was also observed. Since the materials contain very low or no PACs, it was suggested that tumor development may have resulted from chronic skin irritation. CONCAWE, therefore, carried out a program of studies that examined the effect of skin irritation on the tumorigenicity of kerosene (CONCAWE 96/62). In the CONCAWE study, it was found that in the absence of skin irritation kerosene skin tumors did not develop.

Jet Fuel A was included in another study, designed evaluate the role of skin irritation in a skin carcinogenicity of middle distillates. In this study, Jet fuel A was applied either twice weekly or intermittently in which dosing was suspended when marked signs of dermal irritation were noted. When the Jet Fuel A was applied three times weekly 44% of the mice developed tumors and this incidence was reduced to 2% in those animals in which dosing was intermittent. The authors concluded that chronic skin irritation may be a necessary, but not sufficient, condition for skin tumorigenicity (Freeman et al, 1993).

In a review of the available information on the carcinogenicity of middle

## 5. Toxicity

Id Kerosene

Date 9/21/2010

distillate fuels, Nessel (1999) concluded that the tumorigenic activity of the fuels was secondary to skin irritation.

(19) (25) (27) (35) (40) (53) (60)

### 5.8.1 TOXICITY TO FERTILITY

**Type** : Reproduction/developmental study  
**Species** : Rat  
**Sex** : Female  
**Strain** : Sprague-Dawley  
**Route of admin.** : Dermal  
**Exposure period** : 14 days pre mating to day 20 of gestation  
**Frequency of treatm.** : Daily, 7 days/week  
**Premating exposure period**  
    **Male** : 14 days  
    **Female** : 14 days  
**Doses** : 165, 330 & 494 mg/kg/day  
**Control group** : Yes  
**other: NOAEL** : 494 mg/kg bw  
**reproductive/developmental toxicity**  
**Method** : OECD Guide-line 421  
**Year** : 1997  
**GLP** : No data  
**Test substance** : Hydrodesulfurized kerosene

**Method** : The study was performed in accordance with OECD guideline 421 with the addition that males were treated for 8 weeks to improve observation of effects on the reproductive system. Also females were weighed 7 times during gestation rather than 4, and at necropsy, 7 organs in addition to the reproductive organs were weighed.

Ten male Sprague Dawley rats (aged approximately eight week old, weighing 275-285g) and 10 females (same age and weighing 183-187g) were treated dermally with kerosene at concentrations of 0 (sham-treated and vehicle control groups), 20, 40 or 60% (v/v) in mineral oil in a dosing volume of 1 ml/kg. Dosage equivalents were 0, 165, 330 and 494 mg/kg. These doses were selected on the basis of the results of a preliminary 2-week range finding study.

Test material was applied daily to the shorn skin of the animals 7 days/week from 14 days pre mating, during 14 days mating and through 20 days of gestation. Collars were fitted to the animals during the dosing period to prevent ingestion of applied materials. After the final dose, the collars were removed and residual test material was wiped from the skin. Males continued treatment through gestation until final female sacrifice on days 4-6 of lactation.

There were two control groups: the vehicle control was given mineral oil only and in the sham-treated group the animals had been fitted with collars and were stroked with the tip of a syringe, but no material was applied.

During the mating period the test material remained on the animal's backs for 6 hours. Prior to pairing, the test material was removed by wiping. Rats were mated overnight on a 1:1 ratio and were separated the following morning. Collars were then applied prior to the next dose being applied. Females were monitored for evidence that mating had taken place. Pregnancy was determined by the presence of a vaginal plug or sperm in a

vaginal lavage sample. If observed, the female was considered to be at day 0 of gestation. Any female that did not show evidence of mating was placed with the same male the following evening. Any female that did not show evidence of mating at the end of a 2 week mating period was presumed pregnant (gestation day 0 = last day of cohabitation).

Animals were checked twice daily for morbidity and mortality during weekdays but only once daily at weekends. Animals were also observed immediately prior to dosing and after the last animal had been dosed for appearance, behavior and motor activity, respiratory function, central nervous system function, excretory function and biological discharges. Effects of test material on the skin were assessed and scored weekly, using Draize scales for erythema and edema and for chronic deterioration. Males were weighed on the first day of dosing, then weekly and on the day of sacrifice. Females were also weighed on the first day of dosing, then weekly until mating was confirmed and thereafter on gestation days 0, 3, 6, 10, 13, 16 and 20 and on post partum days 0 and 4. Food consumption was also monitored on a similar schedule except through the mating period.

Each presumed-pregnant female was observed daily from gestation day 20 for parturition; evidence of dystocia was noted. The day of delivery was designated postpartum day 0. Maternal behavior and appearance were monitored daily until sacrifice.

Each litter was examined as soon as possible after birth to establish the number and sex of pups, stillbirths, live births and the presence of gross abnormalities. Pups were examined daily for presence of milk in their stomachs. Any pup found dead was examined externally and unusual findings were recorded. The body weight of each viable offspring was individually measured and recorded on post partum days 1 and 4.

Adult females that did not deliver were sacrificed on day 25 of gestation. Dams that delivered and maintained their litters until post partum day 4 were sacrificed with their offspring on post partum days 4-6. All males were sacrificed after the females had been killed. All animals were examined macroscopically for structural anomalies and pathological changes, with emphasis on the reproductive organs. The numbers of implantation sites and corpora lutea of each adult female was recorded. No tissues from offspring were retained.

The liver, kidneys, adrenals, thymus, spleen, brain and heart of all parental animals were weighed. In addition the testes and epididymides of parental males were weighed.

Skin from treated sites, ovaries and testes and epididymides were prepared for histological examination. Pathological evaluation was performed on reproductive organs from all males and pregnant females in both control groups and the high dose group and on treated skin from all groups.

### Statistical evaluation

Quantitative data (body weight and food consumption) were analyzed by parametric methods: analysis of variance (ANOVA) and associated F-test, followed by Dunnett's test for multiple comparisons, provided there was statistical significance in the ANOVA. Maternal reproductive data were evaluated by ANOVA followed by group comparisons using Fisher's exact test. Differences between control and treatment groups were considered statistically significant only if the probability of the differences being due to

## 5. Toxicity

Id Kerosene

Date 9/21/2010

**Remark** : chance was less than 5% ( $P < 0.05$ ).  
 : Two preliminary studies were carried out before the reproductive/developmental toxicity study was conducted.

A dermal irritation study to determine an appropriate dosing regime for the main study

A study of percutaneous absorption to ensure that dilution of test material with mineral oil did not prevent percutaneous absorption of kerosene components.

**Result** : Neither of these studies are summarized here. A summary is only provided for the reproductive/developmental toxicity study itself.  
 : One pregnant mid-dose female died before delivery. No other animals died or were prematurely sacrificed and no clinical signs of toxicity were observed.

Skin irritation among males varied from slight to moderate with increasing dose and was most severe in the high dose group. Mild to moderate skin irritation was observed in females at the highest concentration. At terminal sacrifice, no findings were reported except for those on the skin. Microscopic changes were found in the skin of males in the vehicle control and all kerosene-treated groups. In females changes were only observed in the high dose group animals. The skin findings (macroscopic and microscopic) are shown in the following table.

Parameter	Control	Mineral oil	Kerosene (mg/kg)		
			165	330	494
<b>Males</b>					
No animals	10	10	10	10	10
Max. skin irritation score, sum of means					
Week of max severity	-	2	2	5	5
Mean (SD)	0	1.3(1.2)	2.4(0.7)	2.5(2.0)	3.3(2.1)
Min/max score	0	0/3	1/3	0/7	1/7
Gross necropsy observations					
Crust/scab	1	0	0	0	1
Scaly/dry/flaky	0	0	1	2	3
Histopathological observations					
Acanthosis/hyperkeratosis	2	5	8	7	8
Hyperplasia, sebaceous glands	3	5	5	3	5
Inflammation, dermal	2	1	6	6	7
Necrosis, epidermal, focal	1	0	1	1	5
<b>Females</b>					
No animals	10	6	10	10	10
Max. skin irritation score, sum of means					
Week of max severity	6	7	3	4	4
Mean (SD)	0.2(0.6)	0.7(1.0)	0.4(0.8)	1.1(0.9)	2.3(1.8)
Min/max score	0/2	0/2	0/2	0/2	1/7
Gross necropsy observations					
Crust/scab	0	1	1	0	3
Scaly/dry/flaky	0	0	0	0	0
Histopathological observations					
Acanthosis/hyperkeratosis	3	2	5	5	6
Hyperplasia, sebaceous glands					

## 5. Toxicity

Id Kerosene

Date 9/21/2010

	1	0	0	0	1
Inflammation, dermal	0	1	1	1	4
Necrosis, epidermal, focal	0	0	0	0	0

Body weights were unaffected by treatment. However over the course of the 8 weeks, high dose males gained less weight than the controls (201 g compared to 237g for the controls). Food consumption was unaffected by treatment.

High dose males had a higher mean relative kidney weight than controls (0.76 vs 0.66). This was attributed to the lower mean final body weights of the high dose group. No other organ or organ/body weight changes were recorded.

Parameter	Controls		Kerosene (mg/kg)		
	Sham	Oil	165	330	494
No animals	10	10	10	10	10
Fertility index	100%	90%	90%	80%	100%
Litter with liveborn pups	10	9	9	7 <sup>a</sup>	10
Corpora lutea					
Number	169	151	158	122	172
Mean	16.9	16.8	17.6	17.4	17.2
(SD)	(1.9)	(2.4)	(2.0)	(0.8)	(2.9)
Implantation sites					
Number	163	149	155	18	167
Mean	16.3	16.6	17.2	16.9	16.7
(SD)	(1.9)	(2.4)	(1.8)	(1.3)	(2.8)
Pups delivered					
Total	152	131	147	109	150
Mean	15.2	14.6	16.3	15.6	15.0
(SD)	(2.0)	(2.7)	(2.3)	(2.9)	(2.9)
Liveborn	152	130	143	108	148
Livebirth index	100%	99%	97%	99%	99%
Pups dying					
day 0	3	0	1	1	1
days 1-4	2	4	1	1	9 <sup>bc</sup>
Pups surviving					
4 days	147	126	141	106	138
Viability index	97	97	99	98	93 <sup>c</sup>
Pup weight/litter (g)					
day 1 mean	6.9	6.8	7.0	7.0	6.7
day 4 mean	9.9	9.6	10.1	9.9	9.8

<sup>a</sup> one dam died on gestation day 21; the cause of death was unrelated to treatment

<sup>b</sup> significantly different from control (P<0.05)

<sup>c</sup> One dam had a malfunctioning water bottle; when 4 dead pups from this litter are excluded from the analysis, no significant difference from control was found.

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

**Test substance** : No test-material-related microscopic changes were observed in the testes or epididymides of adult male rats or in the ovaries of adult female rats. The chemical composition of the sample of hydrosulfurized kerosene was determined by ASTM method D 1319-1 and the results are tabulated below.

<b>Component</b>	<b>Weight %</b>
Nonaromatics	80.27
Saturates	78.61
Olefins	1.66
Aromatics	19.72
<3-ring PAC	>19.72
3-7 -ring PAC	<0.01

**Reliability** : (1) valid without restriction

(64)

### 5.8.2 DEVELOPMENTAL TOXICITY/TERATOGENICITY

**Species** : Rat  
**Sex** : Female  
**Strain** : Sprague-Dawley  
**Route of admin.** : Inhalation  
**Exposure period** : Six hours each day  
**Frequency of treatm.** : Daily  
**Duration of test** : Gestation days 6 through 15  
**Doses** : Nominal: 100 & 400 ppm, Actual: 106 & 364 ppm  
**Control group** : Yes  
**NOAEL maternal tox.** : 364 ppm  
**NOAEL teratogen.** : 364 ppm  
**Year** : 1979  
**GLP** : Yes  
**Test substance** : As prescribed by 1.1 - 1.4

**Method** : Groups of 20 female presumed-pregnant rats aged 12 weeks were exposed to nominal concentrations of either 100 or 400 ppm kerosene vapor. Exposures were accomplished in chambers for 6 hours each day on days 6 through 15 of gestation. A control group of 20 presumed-pregnant rats of the same age served as controls and were placed in chambers and exposed to room air only. The rats were weighed on days 0, 6, 15 and 20 of gestation and food consumption was measured throughout the study. The animals were also observed daily for clinical signs.

On day 20 of gestation, the animals were anesthetized with chloroform and the visceral and thoracic organs were examined. The uterus was removed and opened and the number of implantation sites and their placement in the uterine horns were recorded. Live and dead fetuses and resorption sites were also recorded and the fetuses were removed, examined externally for abnormalities and weighed.

One third of the fetuses were fixed in Bouin's fluid and were examined subsequently for changes in the soft tissues of the head, thoracic and visceral organs. The remaining fetuses were stained with Alizarin red S and examined subsequently for skeletal abnormalities.

## 5. Toxicity

Id Kerosene

Date 9/21/2010

### Result

#### Statistical analysis

Analysis of the data was performed using the litter as a basic sampling unit. Dunnett's t-test was used to determine statistical significance ( $P < 0.05$ ) with regard to differences between means with near-normal distribution (body weights and food consumption of dams, mean pup weight based on litter averages). Ratios (nidation index and implantation/corpora lutea ratio) were analyzed with a 2 x 2 contingency table with Yates' correction. Wilcoxon Rank Sum was used for discontinuous parameters as measured by the number of abnormal fetuses within a litter.

: There were no compound-related deaths or clinical signs throughout the study.

At necropsy two animals in the 100 ppm group had lung mottling, but this was considered to be an incidental finding.

There were no significant differences in either body weight or food consumption data.

The following is a summary of reproductive data base on observations of the uterine contents on day 20 of gestation

Observation	Historical control*	Dose (ppm)		
		0	100	400
Nidation index (females with implantations/Bred)	55/61	19/20	18/20	18/20
Females dying prior to Cesarean section	0	0	0	0
Live litters	54	19	18	18
Implantation sites (left horn/right horn)	301/363	110/118	106/138	126/114
Resorptions				
Total	43	18	21	12
Litters	24	11	10	7
Dead fetuses				
Total	0	0	0	0
Litters	0	0	0	0
Mean live litter size (fetuses)	11.3	11.1	12.4	12.7
Average fetal weight (g)	4.1	4.3	4.1	3.9

\* Based on 54 litters

Examination of offspring at delivery did not reveal any treatment-related abnormalities.

Examination of Bouin's fixed specimens did not reveal any treatment-related abnormalities.

The sex ratio was also unaffected by treatment.

Skeletal examinations revealed the following:

Dose group	No. fetuses examined	No. fetuses normal	Fetuses common* changes	with unusual skeletal variations
0	140 (19)	60	72 (16)	8 (4)
106.4	150 (18)	69	72 (15)	9 (4)
364	154 (18)	62	84 (18)	8 (5)

\* Fetuses with commonly-encountered changes only

The authors comment that most of the changes, while not strictly normal, are frequently observed in 20-day-old rat fetuses of this strain and source in their laboratory.



## 5. Toxicity

Id Kerosene

Date 9/21/2010

**Test condition** : The changes were not malformations but were mostly related to retarded bone ossification. Neither the frequency nor the character of the changes indicated an adverse effect on fetal growth and development or a teratogenic potential.  
: A vapor of the test material was generated by metering it into a warmed flask and passing compressed air through the flask. This concentrated vapor was further diluted with room air as it entered the chamber. The chamber concentrations were monitored hourly throughout the exposure.

Nominal and mean actual chamber concentrations are as shown:

<b>Chamber concentration (ppm)</b>	
<b>Nominal</b>	<b>Actual</b>
0	0
100	106.4 ± 10.23
400	364 ± 37.53

**Reliability** : (1) valid without restriction

(7)

**Species** : Rat  
**Sex** : Female  
**Strain** : Sprague-Dawley  
**Route of admin.** : Inhalation  
**Exposure period** : Six hours each day  
**Frequency of treatm.** : Daily  
**Duration of test** : Days 6 through 15 of gestation  
**Doses** : Nominal: 100 & 400 ppm, Actual: 102.5 & 394.7 ppm  
**Control group** : Yes  
**NOAEL maternal tox.** : 394.7 ppm  
**NOAEL teratogen.** : 394.7 ppm  
**Year** : 1979  
**GLP** : Yes  
**Test substance** : Jet fuel A

**Method** : Groups of 20 female presumed-pregnant rats aged 12 weeks were exposed to Jet Fuel A at nominal concentrations of either 100 or 400 ppm vapor. Exposures were accomplished in chambers for 6 hours each day on days 6 through 15 of gestation. A control group of 20 presumed-pregnant rats of the same age served as controls and were placed in chambers and exposed to room air only.  
The rats were weighed on days 0, 6, 15 and 20 of gestation and food consumption was measured throughout the study.  
The animals were also observed daily for clinical signs.

On day 20 of gestation, the animals were anesthetized with chloroform and the visceral and thoracic organs were examined. The uterus was removed and opened and the number of implantation sites and their placement in the uterine horns were recorded. Live and dead fetuses and resorption sites were also recorded and the fetuses were removed, examined externally for abnormalities and weighed.

One third of the fetuses were fixed in Bouin's fluid and were examined subsequently for changes in the soft tissues of the head, thoracic and visceral organs. The remaining fetuses were stained with Alizarin red S and examined subsequently for skeletal abnormalities.

Statistical analysis

## 5. Toxicity

Id Kerosene

Date 9/21/2010

### Result

Analysis of the data was performed using the litter as a basic sampling unit. Dunnett's t-test was used to determine statistical significance ( $P < 0.05$ ) with regard to differences between means with near-normal distribution (body weights and food consumption of dams, mean pup weight based on litter averages). Ratios (nidation index and implantation/corpora lutea ratio) were analyzed with a 2 x 2 contingency table with Yates' correction. Wilcoxon Rank Sum was used for discontinuous parameters as measured by the number of abnormal fetuses within a litter.

: There were no test material-related deaths during the study. Eye irritation (or infection) occurred in a dose-related manner. The irritation consisted of discharge from the eye, swollen eyelids or swollen areas around the eye. The signs of irritation lasted from 2 to 8 days with most animals showing signs for 3 days. Irritation occurred at the following frequency.

Control group	2/20 animals
100 ppm group	7/20 animals
400 ppm group	20/20 animals

Body weights and food consumption were unaffected by treatment.

At necropsy the following findings were made.

Controls 1/20 with mottled lungs  
1/20 with distended intestines  
1/20 with fluid-filled and distended uterus

100 ppm group 2/20 with pale lungs, mottling or dark foci  
One of these animals had a fluid-filled uterus  
1/20 fluid-filled cyst on and around right ovary  
1/20 enlarged, fluid-filled cavernous kidneys

400 ppm group 1/20 granular spleen  
1/20 granular liver  
1/20 mottled lungs and fluid-filled uterus

Observations of the uterine contents did not reveal any treatment-related effects. The data are tabulated below.

Observation	Historical control*	Dose (ppm)		
		0	100	400
Nidation index (females with implantations/Bred)	--	12/20	12/20	16/20
Females dying prior to Cesarean section	--	2**	0	0
Live litters	99%	12	12	16
Implantation sites (left horn/right horn)	46%/54%	64/81	67/88	97/108
Resorptions				
Total	326	8	8	9
Litters	51%	6	4	7
Dead fetuses				
Total	1	0	0	0
Litters	1	0	0	0
Live fetuses/Implantation site	92%	137/145	147/155	197/205***
Mean live litter size (fetuses)	12.2	11.4	12.3	12.3
Average fetal weight (g)	3.6	4.0	3.6	3.7

\* Based on 328 litters

\*\* Misdosed and killed

## 5. Toxicity

Id Kerosene

Date 9/21/2010

\*\*\* Includes set of twins on a single placenta

Examination of offspring at delivery did not reveal any treatment-related abnormalities.

Examination of Bouin's fixed specimens revealed three fetuses from two litters in the 100 ppm group which were judged to be smaller than normal.

Otherwise, there were no other treatment-related abnormalities.

The sex ratio was also unaffected by treatment.

Skeletal examination revealed the following:

Dose group	No fetuses examined	No. fetuses normal	Fetuses common* changes	with unusual skeletal variations
0	90 (12)	44	41 (11)	5 (4)
102.5	98 (12)	62	32 (10)	4 (3)
394.7	131(16)	51	64 (15)	16 (6)

\* Fetuses with commonly-encountered changes only

The authors comment that most of the changes, while not strictly normal, are frequently observed in 20-day-old rat fetuses of this strain and source in their laboratory.

The changes were not malformations but were mostly related to retarded bone ossification. Neither the frequency nor the character of the changes indicated an adverse effect on fetal growth and development or a teratogenic potential.

### Test condition

: The sample of Jet Fuel A was analyzed and the following results were obtained.

Parameter	Method	Result
Flash point	ASTM D 56	135°F
Boiling range	ASTM D 86	325-540 °F
Sulfur		0.036 wt %
Benzene		0.006 vol. %
Hydrocarbon types by mass spectrometry ASTM D2425 (vol. %)		
Saturates		
Paraffins		40.6
Monocycloparaffins		27.3
Dicycloparaffins		10.9
Tricycloparaffins		3.3
Total saturates		82.1
Aromatics		
Alkylbenzenes		10.0
Indans & Tetralins		3.6
Dinaphtheneobenzenes		0.9
C10 Naphthalenes		0.2
C11 Naphthalenes		0.1
Biphenyls, etc.		0.1
Fluorenes, etc.		0
Tricyclic aromatics		0
Total aromatics		17.9

A vapor of the test material was generated by metering it into a warmed flask and passing compressed air through the flask. This concentrated vapor was further diluted with room air as it entered the chamber.

The chamber concentrations were monitored hourly throughout the exposure.

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

Nominal and mean actual chamber concentrations are as shown:

**Chamber concentration (ppm)**

<u>Nominal</u>	<u>Actual</u>
0	0
100	102.5 ± 5.7
400	394.7 ± 19.3

Reliability : (1) valid without restriction

(6)

### 5.9 SPECIFIC INVESTIGATIONS

**Endpoint** : Tumorigenicity  
**Type** : Initiation/Promotion assay  
**Species** : Mouse  
**Sex** : Male  
**Strain** : CD-1  
**Route of admin.** : Dermal  
**Vehicle** : Acetone  
**Control group** : Yes  
**Year** : 1989  
**Test substance** : API 81-07

**Method** : The initiating and promoting activity was determined for nine different petroleum streams, including kerosene (sample API 81-07), in this study. Male CD-1 mice approximately 6 weeks old were used.

#### **Test for initiating activity**

50 µl API 81-07 was applied on five consecutive days to the shorn skin of 30 male mice. After a two week rest period, 50 µl of the promoter PMA (0.1 mg/ml) was applied topically to each animal, twice weekly for 25 weeks. Following the last application of PMA, the dosing was terminated and the mice were sacrificed. A limited gross necropsy was carried out on all mice.

A group of 30 mice treated only with acetone served as the negative controls.

A positive control group were given DMBA. Mice in this group received only one initiating dose (50 µl of 1mg/ml in acetone) on day five of the study. Thereafter, promotion with PMA was the same as for the test groups.

#### **Test for promoting activity**

A single 50 µl dose of DMBA (1mg/ml in acetone) as initiating agent was applied to the shorn skin of a group of 30 male mice. After a two week rest period API 81-07 was applied twice weekly to the mice for 25 weeks.

The negative control group received only acetone as a single dose as the initiating dose. This was followed with 25 weeks of treatment with acetone twice weekly.

A further control group received a single dose of DMBA and following a two week rest these mice were sham handled. This group was included to demonstrate that DMBA had been applied at a sub-threshold dose.

After the completion of the promotion phase all animals were sacrificed and were subjected to a limited gross necropsy.

Throughout the study, the mice were observed daily for morbidity and mortality. Animals were also observed for mass formation throughout the

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

study.  
All mice, whether dying or killed in extremis, were subjected to a limited gross necropsy. This consisted of: examination of application site, brain, heart, lungs, liver, kidneys, spleen, testes, urinary bladder, stomach, duodenum, jejunum, ileum, cecum and colon. Skin of the application site was removed, fixed and processed for histological examination. Other lesions of the internal organs were recorded but no further investigation was undertaken.

**Result** : The results demonstrated that API 81-07 was not a tumor initiating agent but did show tumor promoting activity.

The results of the study are summarized in the following table.

No. mice	Masses observed		Latency (weeks)		
	In life	Histopath	First	Mean	Median mass

### INITIATION ASSAY

API 81-07

30	3	3	25	26.7	27
----	---	---	----	------	----

DMBA control

30	30	30	9	10.6	10
----	----	----	---	------	----

Acetone control

30	3	3	16	18.5	18.5
----	---	---	----	------	------

### PROMOTION ASSAY

API 81-07

acetone

30	0	0	0	0	0
----	---	---	---	---	---

DMBA

30	21	22	14	20.2	19
----	----	----	----	------	----

Sham control

DMBA

29	0	0	0	0	0
----	---	---	---	---	---

**Remark** : In another study, the role of chronic acanthosis on tumor promotion by hydrodesulfurized kerosene (API 81-07) was studied. The authors concluded that their results supported the hypothesis that induction of a lasting, albeit mild, hyperplasia is an essential but not sufficient requirement for tumor promotion. They also concluded that subacute inflammation did not appear to be a significant factor in tumor promotion by the kerosene tested (Skisak, 1991).

(26) (72)

## 5.9 IMMUNOTOXICITY

### TEST SUBSTANCE

Category Chemical : Kerosene/Jet Fuel

Test Substance : Jet Fuel A

## 5. Toxicity

Id Kerosene  
Date 9/21/2010

**Test Substance Purity/Composition and Other Test Substance Comments :**

The test substance met Jet A specifications (ASTM D 1655). The test substance contained 14.9% aromatics and 0.2% naphthalenes, and its 10% boiling point, boiling end point and freezing point were 332°F, 532°F and 49.6°F, respectively.

**Category Chemical Result Type :**

**Unable to Measure or Estimate Justification :**

**METHOD**

**Species:**

Rat

**Other Species:**

**Mammalian Strain:**

CrI:CD (SD) IGS BR

**Gender:**

Female

**Number of Animals per Dose:**

10

**Concentration:**

50% Jet A, 50% Mineral oil

**Dose:**

165, 330, 495 mg/kg

**Year Study Performed :**

2007

**Method/Guideline Followed:**

The study followed the U.S. Environmental Protection Agency Health Effects Test Guidelines for Immunotoxicity (OPPTS 870.7800). The activity of NK cells was assayed as described by Reynolds and Herberman (1981) with modification. Total B cells were enumerated using antibodies directed against the CD45RA surface marker. For total T cells, an antibody to CD5 was used. T subsets were evaluated using antibodies to CD4 for T helper cells and CD8 for cytotoxic T cells. The proliferation of splenocytes in the presence of anti-CD3 antibody was performed as described by Guo et al. (2001).

**GLP:**

Yes

**Vehicle Used:**

Yes

**Vehicle Name:**

Mineral oil

**Other Vehicle Name:**

**Vehicle Amount and Units:**

**Method/Guideline and Test Condition Remarks:**

In a 28-d range-finding study, 4 groups of 3 female rats were administered test substance or test substance/vehicle mixtures at dose levels of 500 mg/kg/d. Other studies on jet fuel and related products indicated that doses higher than

## 5. Toxicity

Id Kerosene

Date 9/21/2010

this would be excessively irritating to the skin. In addition, a group of three female rats served as a control and were sham dosed with an empty syringe and otherwise treated in the same manner as the rats receiving the test substance. Rats were exposed to one of the following: Neat Jet A, Jet A rotated to 4 sites on the back, 50% Jet A in 4:1 acetone:olive oil, or 50% Jet A in mineral oil. Doses were adjusted weekly based on most recent body weights. Clinical observations were made daily throughout the study. Body weights were recorded on d8, d1, at dose initiation (d 0), and on d 7, 14, 21, and 28 for all rats. A full macroscopic postmortem examination was performed on all rats. The treated and untreated skins from all rats were processed, sectioned, stained with hematoxylin and eosin, and examined microscopically.

### TEST RESULTS

Exposure to Jet A when administered topically to rats for a period of 28 d did not alter body weight and food consumption. Furthermore, no effect was observed in either absolute or relative weights for the two major immune organs, the spleen and the thymus.

### Test Results:

The ability of the female animals to produce IgM antibody against sRBC was not affected following 28 d of exposure to Jet A, when evaluated as either specific activity (AFC/106 spleen cells) or total spleen activity (AFC/spleen).

The proliferative response of T cells was not impaired at any dose level following exposure to Jet A. Jet A did not alter the cytotoxic ability of the NK cells at any of the dose levels utilized.

### Results Remarks:

A range-finding study was conducted to evaluate the dermal irritation potential of Jet A when administered topically, unoccluded, to female CrI:CD (SD)IGS BR rats for a period of 28 d by various dosing regimens. Unoccluded topical administration of jet fuel to rats did not produce signs of overt systemic toxicity in any treatment regimen. All animals gained weight over the course of the range-finding study. However, dermal irritation was evident in all treatment groups, but not in the untreated groups. In the neat Jet A group, irritation was severe and animals had to be euthanized on d 5 of exposure due to excessive irritation. Although rotation of the application site relieved irritation somewhat, more serious signs of irritation such as edema and atonia began to appear by the end of the 28-d period. Application of Jet A in a 4:1 mixture of acetone/olive oil (the vehicle of choice for other dermal assays such as the local lymph node assay) produced irritation that was slightly more severe than that observed in the neat Jet A group with the dosing site rotated. Application of 50% Jet A in mineral oil produced the least amount of irritation, with mild to moderate erythema in some animals, desquamation, but no edema. Microscopic examination of tissues from each treatment group was consistent with the degree of irritation observed

## 5. Toxicity

**Id** Kerosene  
**Date** 9/21/2010

(Figure 1). The most common findings were thickening of the epidermis due to hyperplasia, hyperkeratosis, and sebaceous-gland hyperplasia.

### Conclusion:

In contrast to reported immunotoxic effects of jet fuel in mice, dermal exposure of rats to Jet A did not result in alterations in spleen or thymus weights, splenic lymphocyte subpopulations, immunoglobulin (Ig) M antibody-forming cell response to the T-dependent antigen, sheep red blood cells (sRBC), spleen cell proliferative response to anti-CD3 antibody, or natural killer (NK) cell activity. In each of the immunotoxicological assays conducted, the positive control produced the expected results, demonstrating the assay was capable of detecting an effect if one had occurred. Based on the immunological parameters evaluated under the experimental conditions of the study, Jet A did not adversely affect immune responses of female rats.

### RELIABILITY/DATA QUALITY

**Reliability:**

1

**Reliability Remarks:**

**Key Study Sponsor Indicator:**

### REFERENCE

**Reference:**

Cynthia M. Mann, Vanessa L. Peachee, Gary W. Trimmer, Ji-Eun Lee, Lorraine E. Twerdok, and Kimber L. White, Jr Immunotoxicity Evaluation of Jet A Jet Fuel in Female Rats After 28-day Dermal Exposure. *Journal of Toxicology and Environmental Health, Part A*, 71: 495–504, 2008



## 9. References

Id Kerosene  
Date 9/21/2010

- (1) Air Force (1989)  
The installation restoration program toxicology guide. Oak Ridge, TN: Biomedical and Environmental Information Analysis. Vol. 4. [as cited in ATSDR (Agency for Toxic Substances and Disease Registry). 1998. Toxicological Profile for JP-5 and JP-8. U.S. Department of Health and Human Services, Public Health Service, Atlanta, Georgia].
- (2) API (1973)  
Mutagenicity studies of thirteen petroleum fractions  
American Petroleum Institute Report No. 26-60098
- (3) API (1977)  
Mutagenicity evaluation of kerosine  
American Petroleum Institute Med. Res. Publ. 26-60017
- (4) API (1978)  
Estimation of the mutagenicity of hydrocarbon fractions utilizing the bacterial assay procedure with mammalian tissue metabolic activation  
American Petroleum Institute HESD Publ. 26-60103
- (6) API (1979)  
Inhalation/teratology study in rats: Jet Fuel A  
American Petroleum Institute Med. Res. Publ. 27-32173
- (7) API (1979)  
Teratology study in rats: Kerosine  
American Petroleum Institute med. res. Publ. 27-32175
- (8) API (1980)  
Acute toxicity tests of API jetfuel A  
American Petroleum Institute Report No. 27-32815
- (9) API (1980)  
Mutagenicity evaluation of Jet fuel A in the mouse dominant lethal assay  
American Petroleum Institute Med. Res. Publ. 28-31345
- (10) API (1982)  
Acute oral toxicity study in rats  
acute dermal toxicity study in rabbits  
primary dermal irritation study in rabbits  
primary eye irritation study in rabbits  
API 81-07 hydrodeslphurized kerosine (CAS 8008-20-6)  
American Petroleum Institute Med. Res. Publ. 30-31986
- (11) API (1983)  
28-day dermal toxicicty study in the rabbit. Hydrodesulfurized kerosine Sample 81-07  
American Petroleum Institute Med. Res. Publ. 30-32297
- (12) API (1983)  
LC50 Acute inhalation toxicity evaluation of a petroleum derived hydrocarbon in rats.  
Hydrodesulphurized kerosine API Sample 81-07  
American Petroleum Institute Med. Res. Publ. No. 30-32855
- (13) API (1984)  
Dermal sensitization study in guinea pigs, closed patch technique, API sample 81-07  
American Petroleum Institute Med. Res. Publ. 31-31413

## 9. References

Id Kerosene  
Date 9/21/2010

- (14) API (1984)  
Mutagenicity evaluation studies  
in the rat bone marrow cytogenetic assay  
in the mouse lymphoma forward mutation assay  
Hydrodesulfurized kerosine API sample 81-07  
American Petroleum Institute Med. Res. Publ. 32-30240
- (15) API (1985)  
28-day dermal toxicity study of API 83-09 in the rabbit  
American Petroleum Institute report 33-30443
- (16) API (1985)  
Acute oral toxicity study in rats  
acute dermal toxicity study in rabbits  
primary dermal irritation study in rabbits  
primary eye irritation study in rabbits  
dermal sensitization study in guinea pigs  
API 83-09 straight run kerosine (CAS 8008-20-6)  
American Petroleum Institute HESD Res Publ. 32-32858
- (17) API (1985)  
L5178Y TK+/- Mouse lymphoma assay  
API 83-09, Straight run kerosine (CAS 8008-20-6)  
American Petroleum Institute Med. Res. Publ. 32-32745
- (18) API (1985)  
The acute in vivo cytogenetics assay in male and female rats of API 83-09  
American Petroleum Institute Med. Res. Publ. 32-31769
- (19) API (1985)  
The evaluation of the carcinogenicity of certain petroleum fractions  
American Petroleum Institute Med. Res. Publ. 32-30964
- (20) API (1986)  
Four week subchronic inhalation toxicity study in rats  
Final report.  
API 81-07 hydrodesulfurized kerosine (petroleum) (CAS 64742-81-0)  
API 81-09 hydrodesulfurized middle distillate (petroleum) (CAS 64742-80-9)  
API 81-10 hydrodesulfurized middle distillate (petroleum) (CAS 64742-80-9)  
American Petroleum Institute HESD Pub. 33-32724
- (21) API (1987)  
Acute inhalation toxicity of a petroleum derived hydrocarbon in rats.  
API 83-09 Straight run kerosine (CAS 8008-20-6)  
American Petroleum Institute HESD Publ. No. 34-30634
- (22) API (1987)  
Comprehensive Analytical Analysis of API Generic Refinery Streams.  
American Petroleum Institute. Washington, DC.
- (23) API (1988)  
In vivo chromatid exchange assay with API 81-07 Hydrodesulfurized kerosine  
American Petroleum Institute HESD Publ. 36-30043

## 9. References

Id Kerosene  
Date 9/21/2010

- (24) API (1988)  
Sister chromatid exchange assay in Chinese hamster ovary (CHO) cell with API 81-07: hydrodesulfurized kerosine  
American Petroleum Institute HESD Publ. No. 35-32482
- (25) API (1989)  
Lifetime dermal carcinogenesis/chronic toxicity screening bioassay of refinery streams in C3H/HeJ mice  
American Petroleum Institute HESD Publ. No. 36-31364
- (26) API (1989)  
Short-term dermal tumorigenesis study of selected petroleum hydrocarbons in male CD-1 mice.  
Initiation and promotion phases.  
American Petroleum Institute HESD report No. 36-32643
- (27) API (1989)  
Twenty-four month dermal carcinogenesis/chronic toxicity screening bioassay of refinery streams in C3H/HeJ mice (AP-190R). Final report  
American Petroleum Institute Report No. 36-33220
- (28) ASTDR (1998)  
Toxicological profile for JP-5 and JP-8  
Agency for Toxic Substances and Disease Registry  
US Department of health and human services
- (29) ASTM (1999)  
Standard Test Method for Pour Point of Petroleum Oils. ASTM D97, Volume 05.01,  
American Society for Testing and Materials, West Conshohocken, PA.
- (30) ATSDR (1995)  
Jet Fuels (JP-4 and JP-7).  
ATSDR (Agency for Toxic and Disease Registry) Toxicological Profiles, U.S. Department of Health and Human Services, Public Health Service, Atlanta, Georgia
- (31) Battelle (1997)  
13-Week subchronic dermal study with neurotoxicology evaluations of hydrodesulfurized kerosine in Sprague-Dawley rats  
Final report of Battelle study No. N001450A  
Battelle, Columbus, Ohio
- (32) Blackburn, G. R., Deitch, R. A., Schreiner, C. A. and Mackerer, C. R. (1986)  
Predicting tumorigenicity of petroleum distillation fractions using a modified Salmonella mutagenicity assay  
Cell Biol. Toxicol. Vol. 2. pp 63-84
- (33) Bridie, A.L. and Bos, J. (1971)  
Biological degradation of mineral oil in sea water.  
J Inst Petrol 57(557):270-277.
- (34) Carpenter. C. P., Geary, Jr. D. L., Myers, R. C., Nachreiner, L. J. and King, J. M. (1976)  
Petroleum hydrocarbon toxicity studies  
XI Animal and human response to vapors of deodorized kerosine.  
Tox. and Appl. Pharmacol. Vol. 36, 443-456

## 9. References

Id Kerosene  
Date 9/21/2010

- (35) Clark, C. R., Walter, M. K., Ferguson, P. W. and Katchen, (1988)  
Comparative dermal carcinogenesis of shale and petroleum-derived distillates  
Toxicol. and Ind. Health Vol 4, pp 11-22
- (36) CONCAWE ( 1994)  
Kerosines/jet fuels  
Product dossier No. 94/106  
CONCAWE, Brussels
- (37) CONCAWE (1991)  
Middle distillates - a review of the results of a concawe programme of short-term biological studies  
CONCAWE report No. 91/51
- (38) CONCAWE (1993)  
Middle distillates programme, phase 2: overview of the results of a 13-week skin-painting study  
CONCAWE report No. 93/55
- (39) CONCAWE (1995)  
Acute aquatic toxicity of kerosines: report on CONCAWE test programme  
Report No. 96/55
- (40) CONCAWE (1996)  
Overview of the CONCAWE middle distillate programme  
CONCAWE report No. 96/62
- (41) CONCAWE (2001)  
Environmental Classification of Petroleum Substances - Summary Data and Rationale.  
Report No. 01/54
- (42) DSM (1989)  
Acute eye irritation/corrosion study with cracke distillate in albino rabbits  
Report No. V89.199/290069
- (43) DSM (1990)  
Acute (4-hour) inhalation toxicity study with cracked distillate vapour in rats  
Report No. V89/589
- (44) Exxon Biomedical Sciences, Inc. (1995)  
Test Substance MRD-94-883: Daphnia acute toxicity test. Project No. 188342,  
Exxon Biomedical Sciences, Inc. East Millstone, New Jersey.
- (45) Exxon Biomedical Sciences, Inc. (1995)  
Test Substance MRD-94-885: Daphnia acute toxicity test. Project No. 188542,  
Exxon Biomedical Sciences, Inc. East Millstone, New Jersey.
- (46) Exxon Biomedical Sciences, Inc. (1995)  
Test Substance MRD-94-884: Daphnia acute toxicity test. Project No. 188442A,  
Exxon Biomedical Sciences, Inc. East Millstone, New Jersey.
- (47) Exxon Biomedical Sciences Inc. (1996)  
Two year dermal carcinogenicity study of middle distillates in mice  
Final report of project No. 117811A, December 24, 1996

## 9. References

Id Kerosene  
Date 9/21/2010

- (48) Exxon Biomedical Sciences, Inc. (1995)  
Test Substance MRD-94-883: Fish, Acute Toxicity Test, Rainbow Trout - Renewal.  
Project No. 188358,  
Exxon Biomedical Sciences, Inc. East Millstone, New Jersey.
- (49) Exxon Biomedical Sciences, Inc. (1995)  
Test Substance MRD-94-884: Alga, Growth Inhibition Test. Project No. 188467A.  
Exxon Biomedical Sciences, Inc., East Millstone, New Jersey.
- (50) Exxon Biomedical Sciences, Inc. (1995)  
Test Substance MRD-94-884: Fish, Acute Toxicity Test, Rainbow Trout - Renewal.  
Project No. 188458A,  
Exxon Biomedical Sciences, Inc. East Millstone, New Jersey.
- (51) Exxon Biomedical Sciences, Inc. (1995)  
Test Substance MRD-94-885: Alga, Growth Inhibition Test. Project No. 188567.  
Exxon Biomedical Sciences, Inc., East Millstone, New Jersey.
- (52) Exxon Biomedical Sciences, Inc. (1995)  
Test Substance MRD-94-885: Fish, Acute Toxicity Test, Rainbow Trout - Renewal.  
Project No. 188558,  
Exxon Biomedical Sciences, Inc. East Millstone, New Jersey.
- (53) Freeman, J. J., Federici, T. M. and McKee, R. H. (1993)  
Evaluation of the contribution of chronic skin irritation and selected compositional  
parameters to the tumorigenicity of petroleum middle distillates in mouse skin.  
Toxicology. Vol 81, pp 103-112
- (54) Harris, J.C. (1982)  
Rate of Hydrolysis.  
In Handbook of Chemical Property Estimation Methods.  
Lyman, Reehl and Rosenblatt, eds. McGraw-Hill Book Co., New York.
- (55) IARC (1989)  
IARC Monographs on the evaluation of carcinogenic risks to humans, Volume 45:  
occupational exposures in petroleum refining; crude oil and major petroleum fuels.  
Lyon: International Agency for Research on Cancer
- (56) Jokuty, P., S. Whitar, Z. Wang, M. Fingas, B. Fieldhouse, P. Lambert, and J. Mullin.  
(2002)  
Properties of Crude Oils and Oil Products.  
Environmental Protection Service, Environment Canada, Ottawa, Ontario. Internet Version  
2002 via <http://www.etcentre.org/spills>.
- (57) Mackay, D. (1991)  
Multimedia Environmental Models: The Fugacity Approach. Lewis Publ. CRC Press, Boca  
Raton, Florida.
- (58) Mobil (199)  
Determination of the Aerobic Ready Biodegradability of Nigerian Diesel Fuel Using the  
OECD 301F Manometric Respirometry Test Method.  
Mobil Business Resources Corp., Ecotoxicology Laboratory, Paulsboro, NJ, 1999.

## 9. References

Id Kerosene  
Date 9/21/2010

- (59) Mobil (1999)  
Determination of the Aerobic Ready Biodegradability of MSDW Kerosine Mid Blend Using the OECD 301F Manometric Respirometry Test Method.  
Mobil Business Resources Corp., Ecotoxicology Laboratory, Paulsboro, NJ, 1999.
- (60) Nessel, C. S. (1999)  
A comprehensive evaluation of the carcinogenic potential of middle distillate fuels.  
Dru Chem. Toxicol. Vol 22, pp 165-180
- (61) NTP (1986)  
Toxicology and carcinogenesis studies of marine diesel fuel and JP-5 navy fuel (CAS No. 8008-20-6) in B6C3F1 mice (dermal studies).  
Technical report series No. 310  
Research Triangle Park, NC, US Dept of Health and Human Resources
- (62) Potter, T.L. and K.E. Simmons (1998)  
Total Petroleum Hydrocarbon Criteria Working Group Series. Volume 2: Composition of Petroleum Mixtures.  
Amherst Scientific Publishers, Amherst, Massachusetts. 102 pp.
- (63) Rozkov, A., A. Kaard, and R. Vilu (1998)  
Biodegradation of dissolved jet fuel in chemostat by a mixed bacterial culture isolated from a heavily polluted site.  
Biodegradation. 8:363-369.
- (64) Schreiner, C., Bui, Q., Breglia, R., Burnett, D., Koschier, F., Podhasky, P., Lapadula, L., White, R., Feuston, M., Krueger, A. and Rodriguez, S. (1997)  
Toxicity evaluation of petroleum blending streams: reproductive and developmental effects of hydrodesulfurized kerosine  
J. Tox. and Env. Health Vol 52, pp 211-229
- (65) Shell Research (1991)  
The toxicology of petroleum products: the skin irritancy of odourless kerosine (white spirit)  
External report No. SBER 91-001  
Shell Research, Sittingbourne
- (66) Shell research (1991)  
Toxicology of petrochemicals: the skin irritancy of hydrocracked kerosine  
External report No. SBER 91-004  
Shell Research, Sittingbourne
- (67) Shell research (1991)  
Toxicology of petroleum products: the skin irritancy of kerosine SG  
External report No. SBER 91-003  
Shell Research, Sittingbourne
- (68) Shell Research (1991)  
Toxicology of petroleum products: the skin irritancy of AVTUR Jet A1  
External report No. 91-005  
Shell Research, Sittingbourne
- (69) Shell Research Limited. (1995)  
Acute toxicity of water accommodated fractions to Daphnia magna.  
Study No. 6165, Report No. SBGR.94.001.

## 9. References

Id Kerosene  
Date 9/21/2010

- (70) Shell Research Limited. (1995)  
Acute toxicity of water accomodated fractions to Oncorhynchus mykiss.  
Study No. 6165, Report No. SBGR.94.001.
- (71) Shell Research Limited. (1995)  
Acute toxicity of water accomodated fractions to Raphidocelis subcapitata.  
Study No. 6165, Report No. SBGR.94.001.
- (72) Skisak, C. (1991)  
The role of chronic acanthosis and subacute inflammation in tumor promotion in CD-1 mice  
by petroleum middle distillates.  
Tox. and Appl. Pharmacol. Vol 109, pp 399-411
- (73) U.S. EPA (2000)  
EPI (Estimation Programs Interface) Suite, V3.10, Subroutine AOPWIN, V1.90.  
U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics,  
Washington, DC.
- (74) U.S. EPA (2000)  
EPI (Estimation Programs Interface) Suite, V3.10, Subroutine KOWWIN, V1.66.  
U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics,  
Washington, DC.
- (75) U.S. EPA (2001)  
EPI (Estimation Programs Interface) Suite, V3.10. Subroutine WSKOW, V1.40.  
U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics,  
Washington, DC.